

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/274398194>

# The Empirical Status of Psychodynamic Psychotherapy – An Update: Bambi's Alive and Kicking

Article in *Psychotherapy and Psychosomatics* · March 2015

DOI: 10.1159/000376584 · Source: PubMed

---

CITATIONS

26

---

READS

470

4 authors, including:



[Falk Leichsenring](#)

Justus-Liebig-Universität Gießen

226 PUBLICATIONS 5,111 CITATIONS

[SEE PROFILE](#)



[Frank Leweke](#)

Justus-Liebig-Universität Gießen

99 PUBLICATIONS 1,019 CITATIONS

[SEE PROFILE](#)

# The Empirical Status of Psychodynamic Psychotherapy – An Update: Bambi's Alive and Kicking

Falk Leichsenring Frank Leweke Susanne Klein Christiane Steinert

Clinic of Psychosomatics and Psychotherapy, Justus-Liebig University Giessen, Giessen, Germany

## Key Words

Psychodynamic psychotherapy · Empirically supported treatments · Psychotherapy outcome research · Evidence-based medicine

## Abstract

**Background:** The Task Force on Promotion and Dissemination of Psychological Procedures proposed rigorous criteria to define empirically supported psychotherapies. According to these criteria, 2 randomized controlled trials (RCTs) showing efficacy are required for a treatment to be designated as 'efficacious' and 1 RCT for a designation as 'possibly efficacious'. Applying these criteria modified by Chambless and Hollon, this article presents an update on the evidence for psychodynamic therapy (PDT) in specific mental disorders. **Methods:** A systematic search was performed using the criteria by Chambless and Hollon for study selection, as follows: (1) RCT of PDT in adults, (2) use of reliable and valid measures for diagnosis and outcome, (3) use of treatment manuals or manual-like guidelines, (4) adult population treated for specific problems and (5) PDT superior to no treatment, placebo or alternative treatment or equivalent to an established treatment. **Results:** A total of 39 RCTs were included. Following Chambless and Hollon, PDT can presently be designated as efficacious in major depressive disorder (MDD), social anx-

ety disorder, borderline and heterogeneous personality disorders, somatoform pain disorder, and anorexia nervosa. For MDD, this also applies to the combination with pharmacotherapy. PDT can be considered as possibly efficacious in dysthymia, complicated grief, panic disorder, generalized anxiety disorder, and substance abuse/dependence. Evidence is lacking for obsessive-compulsive, posttraumatic stress, bipolar and schizophrenia spectrum disorder(s). **Conclusions:** Evidence has emerged that PDT is efficacious or possibly efficacious in a wide range of common mental disorders. Further research is required for those disorders for which sufficient evidence does not yet exist.

© 2015 S. Karger AG, Basel

## Introduction

Mental disorders are common and pose a challenge to society because they are associated with high costs and considerable impairment [1–3]. Psychotherapy is the first-line treatment option for most of these disorders. Various forms of psychotherapy are available such as cognitive-behavioral therapy (CBT), interpersonal therapy or psychodynamic therapy (PDT). PDT is frequently used in clinical practice [4, 5]. Earlier reviews, however, reported limited evidence for PDT in specific mental dis-

orders [6, 7]. For this reason, the Task Force on Promotion and Dissemination of Psychological Procedures concluded in 1993 that ‘... it is critical that more efficacy evidence on the outcome of psychodynamic therapies for specific disorders be obtained if this clinically verified treatment is to survive in today’s market’ [6, p. 2]. Applying the rigorous criteria proposed by the Task Force and later modified by Chambless and Hollon [8], Connolly Gibbons et al. [7] reviewed the available evidence for PDT in 2008. They concluded that PDT should be considered as ‘possibly efficacious’ treatment for panic disorder, borderline personality disorder and opiate dependence (the Task Force used the term ‘probably efficacious’ [6], whereas Chambless and Hollon used the term ‘possibly efficacious’ [8]).

Furthermore, the authors considered the combination of PDT and pharmacotherapy as ‘efficacious’ in the treatment of major depressive disorder (MDD) [7]. PDT was considered as promising for the treatment of alcohol abuse in order to indicate that this designation was based on a randomized controlled trial (RCT) which did not fulfil all criteria by Chambless and Hollon (no treatment manual was used) [8]. Several reviews on PDT were published in the meantime [9–13] but did not strictly apply the rigorous criteria by Chambless and Hollon [8]. Thus, more than 6 years after the 2008 review by Connolly Gibbons et al. [7], it seems to be timely to update the review on the evidence of PDT in specific mental disorders.

### **Evidence-Based Medicine and Empirically Supported Treatments**

Several proposals have been made to grade the available evidence of both medical and psychotherapeutic treatments [8, 14–16]. All available proposals regard RCTs as the ‘gold standard’ for the demonstration that a treatment is effective. A review and critical discussion of 20 years of evidence-based medicine was recently given by Seshia and Young [17, 18]. The authors discussed the positive impact of evidence-based medicine but also its limitations. RCTs, for example, although regarded as the gold standard, have their limitations. By applying a high degree of experimental control, RCTs emphasize internal validity at the possible expense of external validity, i.e. generalizability to patients and treatments in clinical practice. They also do not usually provide information on long-term effects, adverse events or effectiveness, i.e. how the treatment in question would have performed in the ‘real world’ [18]. Information about adverse events, for

example, often (but not necessarily) relies on observational studies [19]. For psychotherapy, the most rigorous criteria for efficacy were proposed by Chambless and Hollon [8]. For a designation as efficacious, at least 2 RCTs, controlled single-case experiments or equivalent time-sampling designs, carried out in independent research settings are regarded as necessary, in which the respective treatment is superior to no treatment, placebo or alternative treatments or equivalent to an already established treatment. For the latter (equivalence), Chambless and Hollon regarded a sufficient power to detect moderate differences as necessary. In addition, the use of a treatment manual or logical equivalent, reliable and valid procedures to assess diagnosis and outcome, and an appropriate data analysis are required. If there is conflicting evidence, the preponderance of the data must support the efficacy of the treatment. For a designation of possibly efficacious, 1 RCT (sample size of 3 or more in the case of single-case experiments) suffices in the absence of conflicting evidence [8].

The various types of comparison conditions that may be used in an RCT imply that a treatment may be tested more or less strictly. Thus, they are associated with different levels of evidence [20]. The least strict test of a treatment in question by an RCT is achieved by a comparison with a waiting-list condition since it does not control for the unspecific factors of psychotherapy (e.g. attention). A more stringent test is achieved by comparison with a placebo condition, a treatment as usual (TAU) or an alternative treatment, since unspecific factors are controlled for. The most stringent test of efficacy is achieved by comparison with rival treatments, thus controlling for specific and unspecific therapeutic factors [8, p. 8]. According to Chambless and Hollon [8], a treatment can be considered as efficacious and specific if it has been shown to be superior to placebo or to an alternative bona fide treatment in at least two independent research settings.

In order to update the empirical status of PDT with regard to the issues raised above, a review was carried out focusing on RCTs that fulfil the criteria by Chambless and Hollon.

### **Methods**

#### *Definition of PDT*

PDT operates on an interpretive-supportive continuum [21, 22]. Interpretive interventions enhance the patient’s insight about repetitive conflicts sustaining his or her problems [21, 23]. The prototypic insight-enhancing intervention is an interpretation by which unconscious wishes, impulses or defense mechanisms are

made conscious. Supportive interventions aim to strengthen abilities ('ego functions') that are temporarily not accessible to a patient due to acute stress (e.g. traumatic events) or that have not been sufficiently developed (e.g. impulse control in borderline personality disorder). Thus, supportive interventions maintain or build ego functions [22]. Supportive interventions include, for example, fostering a therapeutic alliance, setting goals or strengthening ego functions such as reality testing or impulse control [21]. The use of more supportive or more interpretive (insight-enhancing) interventions depends on the patient's needs. The more severely disturbed a patient is, or the more acute his or her problem is, the more supportive and less interpretive interventions are required and vice versa [21, 22]. Borderline patients, as well as healthy subjects in an acute crisis or after a traumatic event, may need more supportive interventions (e.g. stabilization, providing a safe and supportive environment). Thus, a broad spectrum of mental problems and disorders can be treated with PDT, ranging from milder adjustment disorders or stress reactions to severe personality disorders such as borderline personality disorder or psychotic conditions [21].

Modern variants of PDT are manual guided and specifically tailored to the respective disorder [12]. A diagnostic framework to assess both conflicts and structural deficits which is also useful for therapy planning and the application of adequate interventions was provided, for example, by the Operationalized Psychodynamic Diagnosis system [24].

#### *Updating the Criteria by Chambless and Hollon*

Chambless and Hollon [8] proposed to consider two treatments as equivalent in efficacy if the test treatment was not significantly inferior to the established efficacious treatment, and no trend was observable in the data that the established efficacious treatment was superior in a study of 25–30 subjects per condition. This proposal by Chambless and Hollon, however, is associated with several problems. With 25–30 subjects per condition only large differences can be detected with a sufficient power [25]. Referring to a meta-analysis that had reported a median sample size of 12 per treatment [26], Chambless and Hollon regarded their proposal (25–30 participants) as an intermediate step. According to this 1989 meta-analysis, the majority of studies were not sufficiently powered ( $\geq 0.80$ ) for a comparison of active treatments [26]. Nowadays, however, larger studies are possible and available [27]. Another problem in showing equivalence refers to statistical testing. The traditional two-sided test for differences does not consider a margin of equivalence. Thus, using the traditional two-sided test to establish equivalence frequently leads to incorrect conclusions [28–30]. Furthermore, a nonsignificant result only implies that equality cannot be ruled out [28]. A more appropriate test for equivalence is provided by the two one-sided test (TOST) procedure [28, 29]. TOST requires the researcher to define a margin of equivalence ( $-\Delta_E, \Delta_E$ ) [28, 29, 31]. Only if the confidence interval of the empirically found difference is within the equality margin can equivalence be concluded [28, 29]. In contrast to the traditional two-sided test, the hypothesis tested by TOST refers to equality – not to a difference.

For these reasons, we propose to update the criteria of Chambless and Hollon [8] with regard to statistical testing and power in the following way. For showing equivalence, we propose to use a medium effect size as an equivalence margin. In fact, this update corresponds to a proposal that was made by Chambless and Hollon

themselves. In the very same article they required a study testing equivalence to be sufficiently powered to detect a medium effect size. However, this is not compatible with the proposal of 25–30 participants per group they suggested as an intermediate step. This issue remained unclear to a certain extent in the article by Chambless and Hollon, which is why we are recommending updating this criterion. For showing equivalence, we propose the use of TOST nowadays. If a medium effect size of  $d = 0.5$  is accepted as compatible with equivalence, as originally suggested by Chambless and Hollon [8], 70 subjects are required in each of the two groups to show equivalence of means by TOST for two groups at  $\alpha = 0.05$  with a power of 0.80 (nQuery software, www.statsols.com/products/nquery-advisor-nterim). Thus, in this review, we regarded RCTs comparing two active treatments as sufficiently powered if the sample size per group was at least 70.

For the detection of smaller differences between treatments, considerably more subjects may be needed. The magnitude of the difference that is regarded as clinically significant depends on clinical judgments. For serious outcomes such as mortality smaller thresholds may be more appropriate [32, 33].

In showing equivalence with an established treatment another aspect is of note. If a new treatment proves to be as efficacious as a treatment already established in efficacy without also including a test of the established treatment in the same study, an inference is required to conclude that the new treatment is efficacious. We infer from the previous studies in which the established treatment had proved to be efficacious that it was also efficacious in the RCT testing the new treatment.

#### *Inclusion Criteria*

To be included in this review, studies had to fulfil the following criteria, which are consistent with the proposal by Chambless and Hollon [8]: (1) PDT according to the definition above, (2) RCT, (3) reliable and valid measures for diagnosis and outcome, (4) an adult population treated for specific problems, (5) PDT proven to be superior to no treatment, pill (or psychological placebo) or an alternative treatment or equivalent to an established treatment and (6) use of treatment manuals or manual-like guidelines, i.e. clear descriptions of a treatment including the theoretical background, a set of technical recommendations for this specific psychodynamic treatment such as indications, interventions or timing, and detailed case examples as given, for example, by Luborsky [21], Malan [34], Sifneos [35], Davanloo [36], Mann [37], Strupp and Binder [38], or Horowitz and Kaltreider [39].

We collected studies of PDT in adults that were published between 1970 and October 14th, 2014 by use of a computerized search of PubMed and PsycINFO. The following search terms were used: (psychodynamic or dynamic or psychoanalytic\*) and (therapy or psychotherapy or treatment) and (study or studies or trial\*) and (outcome or result\* or effect\* or change\*) and (psychiat\* or mental\* or psycho\*) and (RCT\* or control\* or compar\*). Manual searches of previous review articles, textbooks and reference lists of included studies were also performed. In addition, we communicated with authors and experts in the field. After completing the literature searches, all hits ( $n = 3,042$ ) were saved in EndNote. After the removal of duplicates ( $n = 274$ ), the authors independently screened titles and abstracts of the resulting 2,768 articles according to the selection criteria described above. All potentially relevant articles were then retrieved for full-text review, which resulted in the inclusion of 39 RCTs (table 1).

### *RCTs of PDT in Specific Mental Disorders*

In addition to the 9 studies listed by Connolly Gibbons et al. [7] for specific mental disorders [40–48], 30 more recent RCTs of PDT fulfilling the criteria by Chambless and Hollon [8] were identified. We did not include RCTs of irritable bowel syndrome or functional dyspepsia as they represent functional somatic disorders. Of the 39 RCTs included, 8 were sufficiently powered to test for equivalence to an established treatment (table 1) [49–56].

### *Models of PDT*

In the studies identified, various forms of PDT were applied (table 1). The models by Luborsky [21], Malan [34], Hobson [57], and Shapiro [58] were used most frequently.

### *Evidence for the Efficacy of PDT in Specific Mental Disorders*

The studies of PDT included in this review are presented for the different mental disorders. Table 1 contains all studies that fulfilled the Chambless and Hollon criteria [8]. RCTs of PDT that fulfilled some, but not all, criteria are presented in a web appendix, including the reasons for exclusion (see online suppl. table 1, web appendix; for all online suppl. material, see [www.karger.com/doi/10.1159/000376584](http://www.karger.com/doi/10.1159/000376584)). It is worth listing these studies as they may be included in future meta-analyses.

### *Depressive Disorders*

In their 2008 review, Connolly Gibbons et al. [7] designated PDT in the context of medication usage as efficacious in MDD. This designation was based on 2 RCTs [40, 41]. In addition, PDT was considered as possibly efficacious in the treatment of MDD in geriatric patients based on the RCT by Thompson et al. [42]. Our search identified a total of 12 RCTs of PDT either alone or combined with pharmacotherapy in depressive disorders that fulfil the criteria by Chambless and Hollon [40–42, 49, 50, 53, 59–64]. With 2 exceptions [62, 64], MDD was studied.

**Major Depressive Disorder.** de Jonghe et al. [49] compared PDT alone with the combination of PDT and pharmacotherapy in MDD. No significant differences between the conditions were found with regard to rates of response and remission in observer-rated measures. Another RCT studied short-term PDT, long-term PDT and solution-focused therapy in patients with depressive or anxiety disorders [53]. Results were presented separately for patients with depressive or anxiety disorders. Short-term PDT was superior to long-term PDT (and as efficacious as solution-focused therapy) with regard to recovery from MDD at the 7-month follow-up, which corresponded to the posttherapy assessment in short-term PDT. No formal treatment manual was used in short-term PDT, but the treatments followed the approaches by Malan [34] and Sifneos [35], which were used as a general manual-like guideline (personal communication with Paul Knekt, January 17, 2014). In the study by Driessen et al. [50] PDT was compared with CBT. In this study the same treatment concept was used as in the RCT by de Jonghe et al. [49]. In contrast to the earlier studies comparing PDT with a bona fide treatment listed below (see also online suppl. table 1), the studies by de Jonghe et al. [49], Knekt et al. [53] and Driessen et al. [50] were sufficiently powered to test for equivalence (table 1). This is also true for the earlier RCT by de Jonghe et al. [41] included by Connolly Gibbons et al. [7]. It is of note, however, that none of these studies applied the TOST procedure but used the traditional two-sided test. Only the RCT by Driessen et al. [50] tested for noninferiority. In this RCT, noninferiority of

PDT compared with CBT could not be shown for the primary outcome, i.e. remission rates at posttreatment (21 vs. 24%), but only for continuous measures of depression. From this trial Thase [65, p. 954] concluded: ‘On the basis of these findings, there is no reason to believe that psychodynamic psychotherapy is a less effective treatment of major depressive disorder than CBT.’ According to these results, PDT as a class can now be considered as efficacious in MDD (table 1; online suppl. table 2, web appendix).

**PDT Combined with Pharmacotherapy.** In addition to the RCT by de Jonghe et al. [41] that had already been included by Connolly Gibbons et al. [7], a further trial was identified. In this RCT, PDT combined with pharmacotherapy was compared with supportive therapy also combined with pharmacotherapy in MDD [63]. At the end of treatment, no differences were found. At the end of the 6-month continuation phase (only pharmacotherapy), however, PDT was significantly superior to supportive therapy with regard to remission rates (72 vs. 12.5%).

Consistent with the conclusion drawn by Connolly Gibbons et al. [7], PDT combined with pharmacotherapy can be considered as efficacious in MDD.

**Specific Populations with Depressive Disorders.** In several RCTs, specific populations with depressive disorders were examined (MDD in geriatric patients, postpartum MDD, depressive disorders in patients with breast cancer) [42, 59, 64]. In these studies, PDT was superior to a waiting list [42] or TAU [59, 64]. In another RCT, PDT and brief supportive therapy in minor depressive disorders (dysthymic disorder, depressive disorder not otherwise specified, adjustment disorder with depressed mood) were superior to a waiting-list condition at the end of treatment, with no differences between the treatments [62]. At the 6-month follow-up, PDT was superior to brief supportive therapy. Thus, PDT can be considered as possibly efficacious in these specific populations with depressive disorders.

**PDT through the Internet in MDD.** In 2 RCTs, Internet-guided self-help based on PDT was studied [60, 61]. In the first study, PDT through the Internet was superior to a structured support intervention (psychoeducation and scheduled weekly contacts online) in MDD [61]. Treatment effects were maintained at the 10-month follow-up. In the second study, PDT through the Internet was superior to a waiting-list condition with regard to remission from MDD (no diagnosis of MDD) in patients with depressive and anxiety disorders [60]. Results were evaluated separately for depressive and anxiety disorders. The effects of the treatment group were maintained at the 7-month follow-up. According to these results, PDT through the Internet can be considered as possibly efficacious in MDD (2 RCTs, but research settings were not independent).

Several RCTs of manual-guided PDT in depressive disorders were identified which did not fulfil all inclusion criteria. They either compared PDT with an established treatment without including another control condition (e.g. waiting list, TAU, placebo) but were not sufficiently powered for demonstrating equivalence [66, 67] or the results were not in favor of PDT [68]. These RCTs are listed in online supplementary table 1 (web appendix), including the reasons for exclusion.

### *Pathological Grief*

In 2 RCTs the treatment of prolonged or complicated grief by short-term psychodynamic group therapy was studied (table 1) [69, 70]. In the first study, short-term psychodynamic group therapy was significantly superior to a waiting list [69]. In the second study, interpretive therapy was superior to supportive therapy with

**Table 1.** Randomized controlled studies of PDT in specific mental disorders that fulfil the criteria for efficacy by Chambless and Hollon [8] updated for comparisons with treatments established in efficacy

Study	PDT, n	Comparison group(s)	Concept of PDT	Duration of PDT
<i>MDD</i>				
Cooper et al. [59], 2003	50	CBT: n = 43 Counselling: n = 48 TAU: n = 52	Cramer et al. [154], Stern [155]: <i>PDT</i> : Exploring the mother's representation of and relationship to her infant taking her own attachment history into account	10 sessions
de Jonghe et al. [49], 2004	106	PDT + pharmacotherapy: n = 85	de Jonghe [117]: <i>Psychoanalytic supportive psychotherapy</i> : (a) Focus on actual relationships (b) Supportive attitude (e.g. empathic, accepting, affirmative, active) (c) Systematic use of supportive interventions (e.g. reducing anxiety, reassuring, encouraging, advising, modeling, confronting, clarifying, reframing symptoms as problem-solving attempts) (d) Defenses are respected (e) Interpretation is used cautiously (f) Transference is used, but not interpreted	16 sessions
Driessen et al. [50], 2013	177	CBT: n = 164	de Jonghe [117], de Jonghe et al. [156]: See above	16 sessions
Johansson et al. [61], 2012	46	Structured support: n = 46	Silverberg [157]: <i>Internet-guided self-help</i> : 9 modules: (1) Introduction to the treatment and the concept by Silverberg (2) Techniques on how to discover unconscious patterns (3) Understanding the patterns (4) Techniques to break unhelpful patterns (5) Preventing relapse to old patterns (6) Applying obtained knowledge in working life (7) Applying obtained knowledge in personal relationships (8) Relationship between unconscious patterns and depression (9) Summary and advice for future All treatment modules end with an encouragement for the participants to try out the strategies described in the particular module and write to the therapists about the experiences from this; the therapists give feedback on the clients' experiences and administer the gradual access to the modules	10 weeks
Thompson et al. [42], 1987 <sup>1</sup>	24	Behavior therapy: n = 25 CBT: n = 27 Waiting list: n = 19	Horowitz and Kaltreider [39]: <i>Brief PDT</i> : (a) Establishing a working alliance (b) Focus on central conflicts, developmental problems, defensive styles making subjects vulnerable to this particular stress experience (c) Use of clarification and interpretation (d) Reappraisal of serious life event (e) Revisions of the inner model of self and world (f) Supportive interventions (g) Termination: working through approaching loss of therapist, relating it to stress event (e.g. loss)	16–20 sessions
<i>Dysthymic disorder</i>				
Maina et al. [62], 2005	10	Supportive therapy: n = 10 Waiting list: n = 10	Malan [34, 158]: <i>Brief dynamic therapy</i> : (a) Early phase: definition of a focus (symptoms, conflicts, or crisis) (b) Middle phase: addressing the focus (c) Terminal phase: discussion of termination, review of progress, consolidation of gains Use of interpretation and clarification to achieve insight into repetitive conflicts and trauma underlying and sustaining the patient's problems	15–30 sessions (mean = 19.6)
<i>Depressive disorders in patients with breast cancer</i>				
Beutel et al. [64], 2014	78	TAU: n = 79	Luborsky [21, 159]: <i>Short-term PDT</i> : (a) Focus on CCRT (b) Interpretive and supportive interventions (c) Specific interventions dealing with life-threatening disease	20–25 sessions

**Table 1** (continued)

Study	PDT, n	Comparison group(s)	Concept of PDT	Duration of PDT
<i>PDT combined with pharmacotherapy in MDD</i>				
Burnand et al. [40], 2002 <sup>1</sup>	35	Clomipramine: n = 39	Burnand et al. [40]: <i>PDT:</i> (a) Psychoeducation (b) Emphasis of the value of the therapeutic relationships (c) Empathic listening (d) Fostering emotional expression (e) Insight into maladaptive interpersonal patterns (f) Facilitation of new interpersonal bonds	10 weeks
de Jonghe et al. [41], 2001 <sup>1</sup>	72	Pharmacotherapy: n = 57	de Jonghe [117]: See above	16 sessions
Maina et al. [63], 2007	18	Brief supportive therapy + pharmacotherapy: n = 17	Malan [34, 158]: See above	15–30 sessions
<i>Mixed samples of patients with depressive and/or anxiety disorders</i>				
Bressi et al. [79], 2010	30	TAU: n = 30	Malan [34, 160]: <i>Short-term PDT:</i> Focus on the central conflict associated with the presented symptoms; therapist taking an active role, selectively disregarding information outside the area agreed on	40 sessions (1 year)
Johansson et al. [60], 2013 <sup>2</sup>	50	Waiting list: n = 50	McCullough et al. [161], Frederick [162]: <i>Internet-guided self-help:</i> 8 modules: (1) Introduction to affect phobia model (2) Explanation of the problem (3) Mindfulness practice (4) Defense restructuring (5) Anxiety regulation (6) Affect experiencing (7) Affect expression and self/other restructuring (8) Summary and advice for continued work	10 weeks
Knekt et al. [53], 2008 <sup>2</sup>	Short-term PDT: 101 Long-term PDT: 128	Solution-focused therapy: n = 97	Malan [34], Sifneos [35] (short-term PDT) Gabbard [23] (long-term PDT) <i>Short-term PDT:</i> (a) Focus on intrapsychic and interpersonal conflicts (b) Transference-based (c) Actively creating alliance (d) Use of confrontation, clarification, interpretation	Long-term PDT: 232 sessions Short-term PDT: 18.5 sessions Solution-focused therapy: 9.8 sessions
<i>Complicated grief</i>				
McCallum and Piper [69], 1990	27	Waiting list: n = 27	McCallum and Piper [163]: <i>Short-term psychoanalytically oriented group psychotherapy:</i> (a) Emphasis on an active therapist role (b) Emphasis on interpretation and clarification relative to support and direction (c) Examining relevant here-and-now events in the group, including transference (d) Encouraging patients to contribute to the therapeutic process of other patients	12 sessions
Piper et al. [70], 2001	53	Supportive therapy: n = 54	McCallum et al. [164]: <i>Interpretive group therapy:</i> (a) Active therapist role (b) Focus on interpretation and transference (c) Encouraging patients to explore uncomfortable emotions (d) Withholding of immediate praise and gratification	12 sessions

**Table 1** (continued)

Study	PDT, n	Comparison group(s)	Concept of PDT	Duration of PDT
<i>Social anxiety disorder</i>				
Bögels et al. [54], 2014	22	CBT: n = 27 Waiting list: n = 27	Malan [34]: PDT: focus on two triangles: (1) "Triangle of conflict" (a) Forbidden thoughts, feelings and wishes which (b) Cause anxiety, and (c) Result in defensive mechanisms such as avoidance (2) Triangle of (a) Parents/caretakers, (b) Important others, and (c) The therapist. Treatment: (a) Early phase: definition of a focus (symptoms, conflicts or crisis) (b) Middle phase: addressing the focus by interpretation and clarification (c) Terminal phase: discussion of termination, review of progress, consolidation of gains	36 sessions
Knijnik et al. [71], 2004	15	Credible placebo control group: n = 15	Knijnik et al. [71]: <i>Psychodynamic group therapy</i> : Phase I: two individual interviews to obtain a psychiatric and developmental history and to conceptualize each patient's focus Phase II: 12 group sessions (a) Sessions 1–3 (address group formation, ensuring patients' agreement to focus exclusively on the treatment of generalized social anxiety disorder and focus formulation) (b) Sessions 4–10 (connection between symptoms and conflicts is investigated) (c) Sessions 11–12 (discuss issues related to treatment termination) Active therapist addressing conflicts and transference emerging in the group and similarities in social problems	12 sessions
Leichsenring et al. [52, 73], 2013, 2014	207	Cognitive Therapy: n = 209 Waiting list: n = 79	Luborsky [21], Leichsenring et al. [165]: <i>Short-term PDT</i> : (a) Focus on core conflictual relationship theme (b) Focus on helping alliance (c) Interpretive and supportive interventions (d) Specific interventions dealing with social fears (e.g. patients are encouraged to confront rather than to avoid the situation they fear)	30 sessions
<i>PDT combined with pharmacotherapy in social anxiety disorder</i>				
Knijnik et al. [74, 75], 2008, 2009	29	Pharmacotherapy: n = 29	Knijnik et al. [74]: See above	12 sessions
<i>Generalized anxiety disorder</i>				
Andersson et al. [77], 2012	27	Internet CBT: n = 27 Waiting list: n = 27	Silverberg [157]: See above	8 weeks
<i>Panic disorder</i>				
Milrod et al. [43], 2007 <sup>1</sup>	26	CBT (applied relaxation): n = 23	Milrod et al. [166]: <i>Panic-focused PDT</i> : Phase I: acute panic (1) Exploration of circumstances/feelings surrounding panic onset (2) Exploration of personal meanings of panic symptoms (3) Exploration of feelings/content of panic episodes Phase II: panic vulnerability (1) Addressing the transference (2) Working through – demonstration that the same conflict emerges in many settings Phase III: termination Addressing patient reaction to termination	24 sessions



**Table 1** (continued)

Study	PDT, n	Comparison group(s)	Concept of PDT	Duration of PDT
<i>PDT combined with pharmacotherapy in panic disorder</i> Wiborg and Dahl [47], 1996 <sup>1</sup>	20	PDT combined with clomipramine (n = 20) vs. clomipramine alone (n = 20)	Wiborg and Dahl [47], Davanloo [36], Malan [34], Strupp and Binder [38]: <i>Brief dynamic psychotherapy:</i> (a) Help patient develop insight into the origins and determinants of his or her dysfunctional patterns (b) Acquisition of more adaptive patterns of interpersonal relatedness (c) Focus on identification and working through of dysfunctional patterns as they emerge in the past, present and transference (d) All technical elements of PDT are included (clarification, confrontation, interpretation of resistance, defensive styles and isolated affects)	15 sessions
<i>Somatic symptom disorder with predominant pain (formerly: somatoform pain disorder)</i> Monsen and Monsen [82], 2000	20	TAU/no therapy: n = 20	Monsen and Monsen [167]: <i>Psychodynamic body therapy:</i> (a) The therapist tries to read affect information, and invites the patients to explore their affect experiences (b) The interventions are adapted to how the patients recognize the affects (awareness) and allow themselves to be moved by the affects (tolerance) (c) Exploration of expressive manners and how certain maladaptive experiencing or expressive patterns are linked to other people and to the representations of significant others (e) Bodily interventions (e.g. massage grips and specific exercises) making affects accessible to conscious awareness	33 sessions
Sattel et al. [83], 2012	107	Enhanced medical care: n = 104	Hardy et al. [168]; PISO Working Group [169]: <i>Brief psychodynamic interpersonal psychotherapy:</i> Phase I: Underscoring the legitimacy of bodily complaint and introducing the possibility of links between emotional states and symptoms of pain; bodily relaxation training is introduced and psychoeducation about the emergence of symptoms Phase II: Clarifying the patient's emotions as they relate to symptoms; discussing these links in the context of actual and past relationships Phase III: Termination issues, such as planning additional psychological, social or pharmacological therapeutic measures	12 sessions
<i>Bulimia nervosa</i> Bachar et al. [84], 1999	17	Cognitive therapy: n = 17 Nutritional counselling: n = 10	Bachar [170]: <i>Self-psychological treatment:</i> The unique conceptualization of self, self-object relations and this theory's conceptualization of resistance and defenses constitutes a therapeutic stance which especially fits the therapeutic needs of eating-disordered patients (a) Therapeutic stance of the therapist as a self-object who tries to empathize with the patient from an experience-near position (b) Conceptualization of food as fulfilling self-object needs	46 sessions
<i>Binge-eating disorder</i> Tasca et al. [92], 2006	48	Group CBT: n = 47 Waiting list: n = 40	Tasca et al. [92]: <i>Group psychodynamic interpersonal psychotherapy:</i> Early stage of the group: Focus on understanding patients' cyclical relational patterns and on helping to develop a cohesive working group Middle stage: Therapist challenges patients' cyclical relational patterns expressed in the group interactions Late stage: Focus on loss and separation as universal stressors, and new cyclical relational patterns and accompanying introjections are reinforced Diet, weight-related issues, and dysfunctional cognitions specific to dietary restraint are not directly addressed	16 sessions

**Table 1** (continued)

Study	PDT, n	Comparison group(s)	Concept of PDT	Duration of PDT
<i>Anorexia nervosa</i> Dare et al. [91], 2001	21	Cognitive-analytic therapy: n = 22 Family therapy: n = 22 Routine treatment: n = 19	Dare [171], Malan [34]: <i>Focal analytic psychotherapy</i> : (a) Time-limited standardized analytic therapy (b) Non-directive (c) No advice about eating behavior or other problems of symptom management Therapist addresses: (a) The unconscious and conscious meanings of the symptom (b) The effects of the symptom on current relationships (c) Manifestation of those influences in the relationship with the therapist (focus on transference)	24.9 sessions
Zipfel et al. [55], 2013	80	Enhanced CBT: n = 80 Optimized TAU: n = 82	Schauenburg et al. [172]: <i>Focal PDT</i> : Identification of psychodynamic foci with a standardized, operationalized, psychodynamic diagnostic interview Three treatment phases: (1) Focus on therapeutic alliance, proanorectic behavior and ego-syntonic beliefs (attitudes and behavior viewed as acceptable) and self-esteem (2) Main focus on relevant relationships and the association with eating (anorectic) behavior (3) Transfer to everyday life, anticipation of treatment termination, and parting Before every treatment session, an independent assessor measured every patient's weight and reported it to his or her therapist	PDT: 39.9 sessions Enhanced CBT: 44.8 sessions Optimized TAU: 50.8 sessions
<i>Substance-related disorders: opiate dependence</i>				
Woody et al. [45, 93], 1983, 1990 <sup>1</sup>	31	Drug counselling: n = 35 CBT + drug counselling: n = 34	Luborsky [21]: <i>Analytically oriented focal therapy</i> : (a) Focus on core conflictual relationship theme (b) Interpretive and supportive interventions (c) Special attention to themes involved in the drug dependence (the role of drugs in relation to problem feelings and behaviors, the meanings that the patient attaches to the drug dependence and how problems may be solved without recourse to drugs)	12 sessions
Woody et al. [46], 1995 <sup>1</sup>	57	Drug counselling: n = 27	Luborsky [21]: See above	26 sessions
<i>Borderline personality disorder</i>				
Bateman and Fonagy [98], 2009	71	Structured clinical management: n = 63	Bateman and Fonagy [173]: <i>Mentalization-based therapy</i> : Focus on increasing the patient's capacity for mentalization and stabilizing the sense of self; mentalization-based therapy requires greater activity, openness and more collaboration on the part of the therapist than in the classical analytical setting Mentalizing stance of the therapist: (a) Self-directed questions (b) Focus on answers that common sense or folk psychology would suggest rather than on unconscious reasons (c) Linking external events to internal states (d) Focus on the psychological process in the here and now (e) Manifest affect is targeted, identified and explored (f) Interpretation of conflict is minimized (g) Continuity between sessions is established (h) Cautious use of transference which is not seen as primary vehicle for change (i) Focus on simple interchanges instead of describing complex mental states	18 months

**Table 1** (continued)

Study	PDT, n	Comparison group(s)	Concept of PDT	Duration of PDT
Clarkin et al. [44], 2007 Levy et al. [95], 2006 <sup>1,3</sup>	30	Dialectical behavioral therapy: n = 30 Supportive therapy: n = 30	Clarkin et al. [99]: <i>Transference-focused psychotherapy</i> : (a) Treatment contract (e.g. duration and payment, potential threats to the treatment specific to each patient, e.g. suicide attempts, drug misuse or anorectic behavior) (b) Integration of internalized experiences of dysfunctional early relationships by focusing on relationship between the individual and the therapist ('transference relationship') (c) Offering a hierarchy of thematic priorities to be used in every session Limits are actively set by the therapist, if a patients' behavior threatens their life, others' lives or the continuation of therapy (d) Phenomenologically, treatment aims at the reduction of impulsivity (e.g. aggression directed towards self or others, substance misuse, eating disorder), mood stabilization and the improvement of interpersonal relationships as well as occupational functioning	12 months
Doering et al. [96], 2010	43	Treatment by experienced community therapists: n = 29	Clarkin et al. [99]: See above	1 year
Gregory et al. [97], 2008	15	TAU: n = 15	Gregory [unpubl. manuscript]; Gregory and Remen [174]: <i>Dynamic deconstructive psychotherapy</i> : (a) Fostering verbalization of affects and of recent interpersonal experiences into simple narratives (b) Helping the patient to integrate polarized attributions toward self and other (e.g. by alternative perspectives or attributions) (c) Focusing on moment-by-moment affective responses of both the patient and therapist (d) Problematic behaviors, including alcohol misuse, are viewed as maladaptive efforts to self-soothe in the absence of verbal/symbolic and relational capacities (e) Therapist encourages verbalization of recent episodes of problematic behaviors within the context of interpersonal narratives (f) Nonjudgmental and nondirective stance	24.9 sessions
<i>Heterogeneous personality disorders</i>				
Abbass et al. [108], 2008	14	Minimal contact: n = 13	Davanloo [36]: <i>Intensive short-term dynamic psychotherapy</i> : (a) Encouraging the awareness and experience of feelings while (b) Clarifying and challenging defenses in collaboration with the patient in order to rapidly help the patient to experience unconscious emotions and work on mobilized transference feelings	27.7 sessions (mean)
Vinnars et al. [56], 2005	80	Community-delivered PDT: n = 76	Luborsky [21]: See above	40 sessions
Winston et al. [48], 1994 <sup>1</sup>	25	Brief adaptive psychotherapy: n = 30; Waiting list: n = 26	Davanloo [36]: See above	40 weeks (mean = 40.3 sessions)
<i>High utilizers of psychiatric services</i>				
Guthrie et al. [110], 1999	55	TAU: n = 55	Hobson [57], Shapiro and Firth [58], Guthrie et al. [175]: <i>Psychodynamic interpersonal therapy</i> : (a) Assumption that the patient's problems arise from or are exacerbated by disturbances of significant personal relationships (b) A tentative, encouraging, supportive approach by the therapist, who seeks to develop deeper understanding with the patient through negotiation, exploration of feelings and metaphor (c) Linkage of the patient's distress to specific interpersonal problems (d) Use of the therapeutic relationship to address problems and test out solutions in the 'here and now'	8 sessions

**Table 1** (continued)

Study	PDT, n	Comparison group(s)	Concept of PDT	Duration of PDT
<i>Relationship distress: marital therapy</i>				
Snyder and Wills [112], 1989 Snyder et al. [113], 1991	30	Behavioral marital therapy: n = 29, Waiting list: n = 20	Nadelson and Paolino [176], Snyder and Wills [112]: <i>Insight-oriented marital therapy:</i> (a) Clarification and interpretation of intra- and interpersonal conflicts (b) Instruction in listening and empathy if necessary (c) Modification of grossly destructive communication patterns No systematic training of communication skills through active rehearsal and behavior-shaping procedures	Up to 25 sessions

<sup>1</sup> Studies fulfilling the criteria by Chambless and Hollon [8] listed by Connolly Gibbons et al. [7].

<sup>2</sup> The outcome was evaluated separately for depressive and anxiety disorders; only results of short-term PDT were included in this review; as for long-term PDT, no manuals were used.

<sup>3</sup> Transference-focused therapy was superior to dialectical behavioral therapy and supportive therapy with regard to measures of attachment and self-reflective functioning [95].

regard to patients achieving reliable or clinically significant change [70]. The treatments can be considered as possibly efficacious (2 RCTs, but research settings were not independent).

#### Social Phobia

For social phobia, we identified 4 RCTs of PDT fulfilling the inclusion criteria. In the first study, short-term psychodynamic group treatment for generalized social phobia was superior to a credible placebo control [71]. In a large-scale multicenter RCT, the efficacy of PDT and cognitive therapy in the treatment of social phobia was studied [52, 72]. Both treatments were significantly superior to the waiting list. Thus, this trial provides evidence that PDT is effective in the treatment of social phobia according to the criteria proposed by Chambless and Hollon [8]. There were no differences between PDT and cognitive therapy with regard to response rates for social phobia (52 vs. 60%) and reduction of depression. Statistically significant differences in favor of cognitive therapy, however, were found with regard to remission rates (36 vs. 26%), self-reported symptoms of social phobia and reduction of interpersonal problems. Differences in terms of between-group effect sizes were small and below the a priori set threshold for clinical significance [52]. In the follow-up study, neither statistically nor clinically significant differences were found between PDT and cognitive therapy [52, 73]. A recent RCT reported PDT to be superior to a waiting list and no differences in outcome were found in comparison with CBT [54]. The authors reported a sufficient power to detect moderate differences [54, p. 370]. Thus, the study seems to provide evidence for equivalent outcome for PDT and CBT. In another RCT, PDT combined with pharmacotherapy was superior to pharmacotherapy alone [74, 75]. Accordingly, PDT can be designated as efficacious in social anxiety disorder and as possibly efficacious in the context of medication use.

#### Panic Disorder

Connolly Gibbons et al. [7] identified 2 RCTs of PDT in anxiety disorders which fulfilled the criteria by Chambless and Hollon [8]. In the study by Milrod et al. [43] PDT was more successful than applied relaxation in panic disorder. In another study, combining pharmacotherapy with PDT was shown to produce significantly

lower relapse rates than pharmacotherapy alone [47]. No additional RCTs fulfilling the inclusion criteria were identified by our search. In a recent RCT, no differences in outcome between PDT and CBT in panic disorder were found [76]. Remission rates at termination and follow-up were 44 and 50% for PDT and 61 and 56% for CBT. However, with 36 and 18 patients the study was not sufficiently powered to demonstrate equivalence (online suppl. table 1, web appendix).

#### Generalized Anxiety Disorder

In generalized anxiety disorder (GAD), 1 RCT fulfilled the inclusion criteria [77]. In a study of Internet-based therapy, there were no differences in outcome between PDT, CBT and waiting list at the end of treatment in the primary outcome measure (Penn State Worry Questionnaire) [77]. With regard to remission rates (no diagnosis of GAD), however, significant differences between the study groups were found, with rates of 54.5, 35 and 16% for PDT, CBT and waiting list, respectively, at the end of treatment. According to these results, PDT delivered over the Internet seems to be as efficacious as Internet-delivered CBT in GAD – the differences in remission were in favor of PDT. Thus, with a sufficient power at most a difference in favor of PDT might have been detected. Accordingly, Internet-delivered PDT can be designated as possibly efficacious in GAD.

In another RCT of GAD, PDT was compared with CBT [78]. No significant differences between PDT and CBT were found with regard to the primary outcome measure. However, in several secondary outcome measures, CBT was superior, both at the end of therapy and at the 6-month follow-up. Other differences may exist that were not detected due to the limited sample size and power. With 28 and 29 patients, the study was not sufficiently powered to demonstrate equivalence (online suppl. table 1, web appendix).

#### Anxiety Disorders

In the study by Knekt et al. [53] mentioned above, short-term PDT was superior to long-term PDT (and as efficacious as solution-focused therapy) with regard to recovery from anxiety disorders at the 7-month follow-up, which corresponded to the post-therapy assessment in short-term PDT.

### Mixed Samples of Depressive and Anxiety Disorders

In an RCT, PDT was superior to TAU in a sample of patients with depressive or anxiety disorders [79].

### Posttraumatic Stress Disorder

Beyond the study by Brom et al. [80] that was already discussed by Connolly Gibbons et al. [7], no RCTs of PDT in posttraumatic stress disorder were identified.

### Obsessive-Compulsive Disorder

There was only 1 RCT of PDT in obsessive-compulsive disorder. In this study PDT combined with pharmacotherapy in obsessive-compulsive disorder was not superior to pharmacotherapy alone [81].

### Somatic Symptom Disorder with Predominant Pain

In the treatment of patients with chronic pain (somatic symptom disorder with predominant pain, formerly named somatoform pain disorder), PDT was superior to a control condition (no treatment or TAU) in measures of pain, psychiatric symptoms, interpersonal problems, and affect consciousness [82]. The results of PDT remained stable or even improved in the 12-month follow-up. In a recent study, PDT was compared with enhanced medical care in patients with multiple somatoform disorders. The presence of a somatoform pain disorder was required for patient inclusion [83]. No differences were found at the end of treatment. However, at the 9-month follow-up PDT was superior to enhanced medical care with regard to improvements in the physical quality of life of the patients [83].

Thus, PDT can be regarded as efficacious in somatoform pain disorder (somatic symptom disorder with predominant pain according to DSM-5).

### Bulimia Nervosa

Only 1 RCT fulfilled the inclusion criteria (table 1). PDT was significantly superior to both a nutritional counselling group and cognitive therapy [84]. This was true of patients with bulimia nervosa and a mixed sample of patients with bulimia nervosa or anorexia nervosa. In a recent RCT, CBT was superior to PDT [85]. This study was challenged for not having used a bona fide treatment for bulimia nervosa [86]. A reply was given by the authors [87]. Thus, conflicting evidence exists for PDT in bulimia nervosa. Judgment is suspended until further research is available.

In 2 further RCTs comparing PDT with CBT, no differences were found in the primary (disorder-specific) outcome measures (bulimic episodes, self-induced vomiting) [88, 89]. Again, however, the studies were not sufficiently powered to detect moderate differences (online suppl. table 1, web appendix). Apart from this limitation, CBT was superior to PDT in some specific measures.

### Anorexia Nervosa

Here, 3 RCTs were identified that fulfilled the inclusion criteria (table 1). In the first RCT, PDT combined with four sessions of nutritional advice was significantly superior to TAU with regard to weight and BMI changes [90]. In another RCT, PDT, cognitive-analytic therapy, family therapy, and routine treatment were compared [91]. No differences between PDT, cognitive-analytic therapy and family therapy were found. PDT and family therapy were significantly superior to the routine treatment with regard to weight gain. A recent RCT compared manual-guided PDT, en-

hanced CBT and optimized TAU in the treatment of anorexia nervosa [55]. At the end of treatment, significant improvements were found in all treatments, with no differences in the primary outcome measure (BMI). At the 12-month follow-up, however, PDT was significantly superior to optimized TAU with regard to rates of recovery, whereas enhanced CBT was not [55]. Recovery rates were 35 versus 19 versus 13% for PDT, enhanced CBT and optimized TAU, respectively. According to these results, PDT can be designated as efficacious in anorexia nervosa.

### Binge-Eating Disorder

In an RCT of binge-eating disorder, PDT was superior to a waiting list [92]. Thus, PDT can be considered as possibly efficacious in binge-eating disorder. No differences were found between PDT and CBT (e.g. days binged, interpersonal problems) [92]. For the comparison of PDT with CBT, again the statistical power was not sufficient (table 1).

### Substance-Related Disorders

Based on 2 RCTs carried out by the same research team (table 1), Connolly Gibbons et al. [7] considered PDT as a possibly efficacious treatment for opiate addiction [45, 46, 93]. As noted already by Connolly Gibbons et al. [7], another RCT failed to demonstrate efficacy for PDT or CBT in the treatment of cocaine dependence [51].

### Alcohol Abuse

No RCTs were identified that fulfilled the inclusion criteria. Connolly Gibbons et al. [7] considered the results by Sandahl et al. [94] as promising.

### Borderline Personality Disorder

Based on the results by Clarkin et al. [44] and Levy et al. [95], Connolly Gibbons et al. [7] considered transference-focused therapy as possibly efficacious in borderline personality disorder.

We identified 3 more recent RCTs fulfilling the inclusion criteria (table 1) [96–98]. Doering et al. [96] compared transference-focused psychotherapy based on the model of Clarkin and Kernberg [99] with a treatment carried out by experienced community psychotherapists in borderline outpatients [96]. Transference-focused psychotherapy was superior with regard to borderline psychopathology, psychosocial functioning, personality organization, inpatient admission, and dropouts. Another RCT compared PDT ('dynamic deconstructive psychotherapy') with TAU in the treatment of patients with BPD and co-occurring alcohol use disorder [97]. In this study, PDT, but not TAU, achieved significant improvements in outcome measures of parasuicide, alcohol misuse and institutional care [97]. Furthermore, PDT was superior with regard to improvements in borderline psychopathology, depression and social support. No difference was found in dissociation. This was true although TAU participants received higher average treatment intensity. Another RCT found mentalization-based treatment to be superior to manual-driven structured clinical management with regard to the primary (suicidal and self-injurious behaviors, hospitalization) and secondary outcome measures (e.g. depression, general symptom distress, interpersonal functioning) [98]. In another RCT, no treatment manual was used [100]. Giesen-Bloo et al. [101] reported schema-focused therapy to be superior to PDT (transference-focused psychotherapy; online suppl. table 1, web appendix). This study has been challenged with

regard to treatment integrity [102]. A reply was given by the authors [103]. In the case of conflicting evidence, the criteria by Chambless and Hollon [8, p 18] require that the preponderance of evidence must be in favor of the empirically supported treatment [8]. As 2 other RCTs described above found transference-focused psychotherapy to be efficacious [44, 95, 96], the preponderance of evidence is in favor of transference-focused psychotherapy.

In sum, PDT as a group can be considered efficacious and specific in borderline personality disorder according to the criteria proposed by Chambless and Hollon [8]. For transference-focused therapy, 2 RCTs carried out in independent research settings are available which provide evidence that the treatment is efficacious and specific in the treatment of borderline personality disorder [44, 95, 96]. The specific methods of PDT used in the studies by Bateman and Fonagy [98] and Gregory et al. [97] can be considered as possibly efficacious.

#### Cluster C Personality Disorders

No RCTs were identified that fulfilled the inclusion criteria. In an RCT by Svartberg et al. [104] both PDT and CBT yielded significant improvements in symptoms, interpersonal problems and core personality pathology (online suppl. table 1, web appendix). No significant differences were found between PDT and CBT. However, symptom change at follow-up was significant for PDT but not for CBT. Although this rate was almost twice as large for PDT than for CBT, the difference was not statistically significant [104, p 813]. This is probably due to the fact that the study was also not sufficiently powered to detect possible differences ( $N_1 = 25$ ,  $N_2 = 25$ ). This applies to another RCT as well [105]. A further study did not provide evidence for PDT in avoidant personality disorder [106].

#### Heterogeneous Samples of Patients with Personality Disorders

Winston et al. [48] compared PDT with brief adaptive psychotherapy or waiting-list patients in a heterogeneous group of patients with personality disorders. Most of the patients showed a cluster C personality disorder. Patients with paranoid, schizoid, schizotypal, borderline, and narcissistic personality disorders were excluded. In both treatment groups, patients showed significantly more improvements than the patients on the waiting list. No differences in outcome were found between the two forms of psychotherapy. However, the study was not sufficiently powered to detect possible differences between bona fide treatments (online suppl. table 1, web appendix). This is also true for the RCT by Hellerstein et al. [107], who compared PDT to brief supportive therapy in a heterogeneous sample of patients with personality disorders (online suppl. table 1, web appendix). Again, most of the patients showed a cluster C personality disorder. The authors reported similar degrees of improvement both at termination and at the 6-month follow-up. Abbas et al. [108] compared PDT (intensive short-term dynamic psychotherapy) with a minimal contact group in a heterogeneous group of patients with personality disorders. The most common Axis II diagnoses were borderline (44%), obsessive compulsive (37%) and avoidant personality disorder (33%). PDT was significantly superior to the control condition in all primary outcomes. In addition, treatment costs were thrice offset by reductions in medication and disability payments. When control patients were treated, they experienced benefits similar to the initial treatment group. In the long-term follow-up 2 years after the

end of treatment, the whole group maintained their gains and had an 83% reduction of personality disorder diagnoses. In another study, manual-guided PDT was as effective as PDT carried out by community experts [56].

According to these results, PDT following the concept by Davanloo [36] can be considered as efficacious in patients with heterogeneous personality disorders.

#### Deliberate Self-Poisoning

In patients with deliberate self-poisoning, psychodynamic-interpersonal therapy was significantly superior to a TAU condition [109] (online suppl. table 1, web appendix). Unfortunately, the patient sample was not described in terms of mental disorders. PDT can be considered as promising in patients with deliberate self-poisoning.

#### High Utilizers of Psychiatric Services

In another RCT, psychodynamic-interpersonal therapy was superior to a TAU condition in high utilizers of psychiatric services [110]. The sample primarily included patients with depressive and anxiety disorders. PDT can be considered as possibly efficacious in high utilizers of psychiatric services with primarily depressive and anxiety disorders.

#### Bipolar Disorders

No RCT was available for PDT in bipolar disorders.

#### Schizophrenia Spectrum Disorders

No RCT fulfilling the inclusion criteria was available. Promising results were reported by a quasi-experimental study for schizophrenia spectrum disorder [111].

#### Relationship Distress: Marital Therapy

In an RCT, no significant differences were found between psychodynamic and behavioral couple therapy in individual and relationship functioning [112]. Both treatments were superior to a waiting-list control group. Effects were maintained at the 6-month follow-up. At the 4-year follow-up, significantly more couples in the behavioral therapy than in the psychodynamic condition had experienced divorce (38 vs. 3%) [113]. Again, however, the study was sufficiently powered for showing superiority but not equivalence.

## Discussion

In times of evidence-based medicine, it is indispensable that there be sufficient evidence for a treatment before it can be recommended for application in clinical practice. This applies to psychotherapy as well. The 2010 Affordable Care Act in the USA, for example, mandates mental health care, including psychotherapy, as one of the 10 essential health benefits [114].

In his now classic article on the evidence for psychotherapy, Parloff [115, p. 718] concluded in 1982 that 'the prospect that rigorous research evidence will soon provide a credible list of "certified" techniques and proce-

dures for the treatment of specific disorders is poor.’ The article was entitled: ‘Psychotherapy research evidence and reimbursement decisions: Bambi meets Godzilla.’ In 2008, Richard Glass [116], deputy editor of JAMA, commented on a meta-analysis of PDT by asking: ‘Psychodynamic psychotherapy and research evidence: Bambi survives Godzilla?’ [116, p 1587]. According to the review presented here, Bambi seems to be alive and kicking. We carried out an update on the evidence for PDT in specific mental disorders under the requirements of the rigorous criteria proposed by Chambless and Hollon [8]. Compared with the 2008 review by Connolly Gibbons et al. [7], evidence for PDT has considerably increased (table 1, online suppl. tables 1 and 2, web appendix). Whereas Connolly Gibbons et al. found PDT to be possibly efficacious in only a few mental disorders, now evidence is available showing that PDT is efficacious or possibly efficacious in most of the common mental disorders (table 1, online suppl. table 2, web appendix). In MDD, social anxiety disorder, borderline personality disorder, somatoform pain disorder, and anorexia nervosa, PDT can now be considered as efficacious. With regard to specific psychodynamic approaches, the treatments by de Jonghe [117] for MDD, Clarkin et al. [99] for borderline personality disorder and Davanloo [36] for heterogeneous personality disorders can be considered as efficacious. In addition, PDT can now be considered as possibly efficacious in dysthymia, prolonged or complicated grief, GAD (through the Internet), panic disorder, binge-eating disorder, post-traumatic stress disorder, and substance-related disorders. The combination of PDT with pharmacotherapy can be considered as efficacious in MDD and as possibly efficacious in panic disorder and social anxiety disorder. For bulimia nervosa, judgment is suspended. In addition, there is evidence from several RCTs that PDT is efficacious in patients with somatic symptoms [12, 118].

For a treatment indication, three dimensions need to be weighed up – the risk associated with withholding the treatment, the responsiveness to the treatment and vulnerability to the adverse effects of a treatment [19, 119]. General contraindications for PDT are not known; PDT can be adapted to the individual patient’s needs by using either more interpretive or supportive interventions. Thus, as noted above, a broad spectrum of mental disorders can be successfully treated with PDT. However, some contraindications exist for psychoanalytic therapy carried out in the classical couch setting with a high frequency of sessions per week, for example for severely disturbed borderline or psychotic patients. Berk and Parker [120] discussed several risks of high-frequency long-term

psychoanalytic therapy in more detail such as promoting dependency on the therapy and the therapist, focusing on patient history at the expense of current issues and the patient’s active role, or reifying the patient’s doubts about their self-worth and interpersonal skills by a nonresponsive interpersonal style. These issues indeed constitute specific risks of long-term and high-frequency psychoanalytic therapy. They are explicitly regarded as treatment mistakes also in psychoanalysis [121]. Furthermore, the risks associated with long-term treatments are not specific to psychoanalysis since some CBT treatments such as dialectical behavior therapy or schema-focused therapy are long-term as well [44, 101]. It is of note that Berk and Parker [120] only referred to high-frequency long-term psychoanalytic therapy but did not include a discussion of manual-guided PDT. Several RCTs that fulfilled the criteria of Chambless and Hollon [8] (table 1) included data on adverse events or side effects [40, 43, 47, 49, 50, 52, 55, 64, 74, 77, 95, 122]. In these studies, adverse events or side effects were not more frequent in PDT than in the comparison treatments. A total of 3 studies encompassed long-term PDT of more than 25 sessions or of a treatment duration of 12 months [55, 95, 122]. In 2 other studies, significantly less adverse or serious adverse events were found for PDT [47, 64]. For marital therapy, long-term follow-ups reported significantly fewer divorces for psychodynamic marital therapy compared with cognitive-behavioral marital therapy (3 vs. 38%) [113]. Hence, there is no reason to believe that manual-guided PDT is associated with more adverse events or side effects than other bona fide treatments. In general, future studies of psychotherapy should carefully monitor and report adverse events or side effects.

In only a few of the RCTs fulfilling the criteria by Chambless and Hollon [8] included here (table 1) were differences in outcome found in direct comparisons with other bona fide treatments such as CBT. In the study by Leichsenring et al. [52] on social anxiety disorder, CBT achieved statistically significant better outcomes in some measures immediately after the end of treatment. However, the differences were small and below the a priori set threshold of clinical significance. Furthermore, in the long-term follow-ups, neither statistically nor clinically significant differences were found [73]. With regard to the treatment of depression, Thase [65, p. 954] concluded from the RCT by Driessen et al. [50] that there is no evidence that PDT is a less effective treatment of MDD than CBT [95]. On the other hand, Milrod et al. [43] found PDT to be superior to applied relaxation in panic disorder. Clarkin et al. [44] reported PDT and dialectical be-

havior therapy to be equally effective. In measures of attachment and reflective functioning, however, PDT was superior to dialectical behavior therapy [95]. In anorexia nervosa, only PDT (not CBT) was superior to optimized TAU in the 12-month long-term follow-up [55].

As noted above, many studies were not sufficiently powered for showing equivalence to a treatment established in efficacy. Studies of CBT do not seem to be more highly powered. In comparisons of psychotherapy with pharmacotherapy in depression, only 2 of 26 studies included at least 70 patients per group [123]. On the other hand, meta-analyses may achieve a higher statistical power than an individual study. In anxiety disorders, a recent meta-analysis found PDT to be as efficacious as other bona fide treatments [124]. For depression, another meta-analysis found little indication that CBT was superior to other forms of psychotherapy such as interpersonal therapy or PDT [125]. In direct comparisons of PDT with CBT, between-group effect sizes were found to be small [126]. Furthermore, the clinical relevance of statistically significant but small differences between treatments is often not clear [18, 127].

If only few and small differences in outcome between PDT and other approaches such as CBT exist, the question arises whether there are patients who benefit from one approach rather than from another. Unfortunately, there is a lack of systematic research on this important question.

A few studies reported some post hoc analyses [92, 128]. Tasca et al. [92], for example, reported that in binge-eating disorder patients with high attachment anxiety did better in psychodynamic group therapy whereas those with low attachment anxiety did better in group CBT. Given the paucity of research, future studies are recommended to formulate and test hypotheses on moderators of treatment outcome [25].

With regard to comparisons with pharmacotherapy, it is of note that no direct comparisons of PDT alone with pharmacotherapy alone are available. CBT was shown to be superior to pharmacotherapy with regard to relapse prevention in depression [129]. As a limitation, it should be noted, however, that this effect was not found in all studies [130]. Furthermore, a sustained response was found only in a limited proportion of patients treated with CBT (37.3%) [129, p. 420]. Nevertheless, for these patients CBT was shown to prevent relapse to a significantly higher degree than pharmacotherapy [129]. A recent meta-analysis found CBT to be superior to discontinued pharmacotherapy with regard to relapse prevention in depression but not superior to continued

pharmacotherapy [125]. For PDT, studies on relapse prevention in depression are not yet available. This is an important area for future research in PDT. As a promising result, remission rates of PDT seem to be stable in follow-up assessments [50, 131], with no differences to CBT [131]. For panic disorder, adding PDT to pharmacotherapy was found to be superior to pharmacotherapy with regard to relapse prevention (75 vs. 20%) [47].

Evidence has emerged that even in studies of manual-guided therapies a considerable overlap exists between interventions [132–135]. Cognitive-behavioral therapists, for example, were found to use interpretations or clarifications as often as psychodynamic therapists [135], and psychodynamic therapists were found to adhere to a CBT prototype to the same degree as they adhered to a psychodynamic prototype [133]. Ablon and Jones [132] reported that the interpersonal therapies in the NIMH Treatment of Depression Collaborative Research Program corresponded most strongly to a prototype of an ideal CBT and that congruence of the interpersonal therapies to the CBT prototype yielded more positive correlations with outcome measures than congruence to the interpersonal prototype. For this reason, Ablon and Jones [132, p. 780] concluded: ‘Brand names of therapy can be misleading.’

More important than the issue of overlap in interventions, however, is the question of which interventions are related to outcome. In several studies, significant relationships between psychodynamic interventions and outcome could be demonstrated [136–141]. These studies provide another type of evidence for PDT, i.e. that outcome is actually related to psychodynamic interventions. Furthermore, there is evidence that changes in psychodynamically relevant mediators such as core conflicts or insight are significantly related to outcome [142, 143].

For some mental disorders, RCTs fulfilling the criteria by Chambless and Hollon [8] providing evidence for the efficacy of PDT do not yet exist. This applies to obsessive-compulsive disorder, posttraumatic stress disorder, bipolar disorders, and schizophrenia. Thus, further research on PDT is required. As a research strategy, it seems to be promising to carry out a second RCT for those methods of PDT for which already 1 RCT provided evidence for efficacy (possibly efficacious treatments). Thus, the online supplementary tables 1 and 2 (web appendix) may provide a guideline for further research. If the second RCT also provides evidence for the efficacy of the specific treatment in question, this treatment can be designated as efficacious. As another approach, developing and testing transdiagnostic and unified treatment protocols may be a



promising strategy. For CBT, several unified protocols are available [144]. For PDT, the first unified protocols for the treatment of anxiety disorders and depressive disorders have recently been published [145, 146]. The unified protocol for anxiety disorders, for example, integrates those treatment elements that were used in RCTs in which PDT proved to be efficacious in the various categories of anxiety disorders [145]. Thus, it represents a transdiagnostic protocol for the treatment of the various categories of anxiety disorders (e.g. GAD, panic disorder, social anxiety disorder) as represented, for example, in DSM-5. Another transdiagnostic approach was presented by Busch et al. [147], who extended the treatment developed for panic disorder to other anxiety disorders. The approach by Busch et al. is also transdiagnostic, but it is not a unified approach as it does not systematically integrate treatment elements of all available evidence-based treatments. At present, evidence for this approach is only available for panic disorder [147]. It may be a strength of PDT that it is transdiagnostic in origin as it does not focus on specific symptoms (e.g. symptoms of panic, generalized anxiety or social phobia) but on the conflicts and structural deficits underlying the symptoms [145].

Evidence for PDT in specific mental disorders usually comes from RCTs in which different concepts and methods of PDT were applied (table 1). Thus, the available evidence for PDT in a specific mental disorder is 'scattered' between the different approaches of PDT applied (table 1). For social anxiety disorder, for example, 3 RCTs were available, each of them using a different form of PDT. It was for this very reason that PDT was judged as only possibly efficacious by Chambless and Hollon [8]. Unified protocols contribute to overcoming this problem [145, 146]. If a unified protocol is used in several RCTs, the evidence is valid for the same treatment concept (the unified protocol).

With regard to the methodological quality of studies included here, this review was based on the rigorous criteria proposed by Chambless and Hollon [8]. Recent meta-analyses also reported that the quality of RCTs of PDT was as good as that of RCTs of CBT [11, 148].

Online supplementary table 1 (web appendix) presents the RCTs that for one reason or another do not fulfil the criteria for efficacy proposed by Chambless and Hollon [8]. In most cases, statistical power was not sufficient to demonstrate equivalence with an already established treatment. Thus, researchers are strongly recommended to ensure a sufficient power if a treatment is to be compared to an established treatment, or they should at least include an additional waiting list or TAU condition in

order to be able to demonstrate efficacy. However, the existing underpowered RCTs are not necessarily useless or unethical [149, 150]. If they are included in a meta-analysis, the statistical power will increase beyond that of the individual study, with the power depending not only on the number of studies and patients but also on the degree of heterogeneity between studies [151].

Our review did not include studies of children and adolescents. A recent meta-analysis provided some evidence that PDT is effective in children and adolescents across a range of common mental disorders [152].

We presented the studies of PDT for the different mental disorders. However, from a psychodynamic perspective, the results of a therapy for a specific mental disorder (e.g. depression, agoraphobia) are influenced by the underlying psychodynamic features (e.g. conflicts, defenses, personality organization), which may vary considerably within one category of psychiatric disorder [153]. These psychodynamic factors may affect treatment outcome and may have a greater impact on outcome than the phenomenological DSM categories [70]. Psychodynamic features as assessed, for example, by operationalized psychodynamic diagnosis [24], may be used in future studies to address the question of which patients benefit from which kind of treatment. They may also be helpful in examining which patients do not respond well to a specific treatment.

In summary, considerable progress has been made concerning the empirical status of PDT, but further research is needed to fill the gaps described above.

## Acknowledgment

We would like to thank Dr. Chambless for some clarifications with regard to the criteria of empirically supported therapies.

## Disclosure Statement

We have no conflicts of interest to declare.

## References

- 1 Kessler RC, Berglund P, Demler O, Jin R, Walters EE: Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the national comorbidity survey replication. *Arch Gen Psychiatry* 2005;62:593–602.
- 2 Kessler RC, Chiu WT, Demler O, Merikangas KR, Walters EE: Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the national comorbidity survey replication. *Arch Gen Psychiatry* 2005;62:617–627.

- 3 Kessler RC, Petukhova M, Sampson NA, Zaslavsky AM, Wittchen HU: Twelve-month and lifetime prevalence and lifetime morbid risk of anxiety and mood disorders in the United States. *Int J Methods Psychiatr Res* 2012;21:169–184.
- 4 Norcross JC, Rogan JD: Psychologists conducting psychotherapy in 2012: current practices and historical trends among Division 29 members. *Psychotherapy (Chic)* 2013;50:490–495.
- 5 Cook JM, Biyanova T, Elhai J, Schnurr PP, Coyne JC: What do psychotherapists really do in practice? An internet study of over 2,000 practitioners. *Psychotherapy* 2010;47:260–267.
- 6 Chambless DL: Task Force on Promotion and Dissemination of Psychological Procedures: A Report Adopted by the Division 12 Board of the American Psychological Association. Oklahoma City, APA, 1993.
- 7 Connolly Gibbons MB, Crits-Christoph P, Hearon B: The empirical status of psychodynamic therapies. *Annu Rev Clin Psychol* 2008;4:93–108.
- 8 Chambless DL, Hollon SD: Defining empirically supported therapies. *J Consult Clin Psychol* 1998;66:7–18.
- 9 Fonagy P, Roth A, Higgitt A: Psychodynamic therapies, evidence-based practice and clinical wisdom. *Bull Menninger Clin* 2005;69:1–58.
- 10 Shedler J: The efficacy of psychodynamic psychotherapy. *Am Psychol* 2010;65:98–109.
- 11 Gerber AJ, Kocsis JH, Milrod BL, Roose SP, Barber JP, Thase ME, Perkins P, Leon AC: A quality-based review of randomized controlled trials of psychodynamic psychotherapy. *Am J Psychiatry* 2011;168:19–28.
- 12 Leichsenring F, Klein S: Evidence for psychodynamic psychotherapy in specific mental disorders: a systematic review. *Psychoanal Psychother* 2014;28:4–32.
- 13 Abbass AA, Kiskey SR, Town JM, Leichsenring F, Driessen E, De Maaat S, Gerber A, Dekker J, Rabung S, Rusalovska S, Crowe E: Short-term psychodynamic psychotherapy for common mental disorders. *Cochrane database Syst Rev* 2014;7:CD004687.
- 14 Canadian Task Force on the Periodic Health Examination: The periodic health examination. *Can Med Assoc J* 1979;121:1193–1254.
- 15 Clarke M, Oxman AD: *Cochrane Reviewers' Handbook 4.1.6* (updated January 2003). Oxford, Update Software, 2003.
- 16 Cook DJ, Guyatt GH, Laupacis A, Sackett DL, Goldberg RJ: Clinical recommendations using levels of evidence for antithrombotic agents. *Chest* 1995;108:227S–230S.
- 17 Seshia SS, Young GB: The evidence-based medicine paradigm: where are we 20 years later? Part 2. *Can J Neurol Sci* 2013;40:475–481.
- 18 Seshia SS, Young GB: The evidence-based medicine paradigm: where are we 20 years later? Part 1. *Can J Neurol Sci* 2013;40:465–474.
- 19 Fava GA: Rational use of antidepressant drugs. *Psychother Psychosom* 2014;83:197–204.
- 20 Gabbard G, Gunderson JG, Fonagy P: The place of psychoanalytic treatments within psychiatry. *Arch Gen Psychiatry* 2002;59:505–510.
- 21 Luborsky L: *Principles of Psychoanalytic Psychotherapy. Manual for Supportive-Expressive Treatment*. New York, Basic Books, 1984.
- 22 Wallerstein R: The psychotherapy research project of the Menninger foundation: an overview. *J Consult Clin Psychol* 1989;57:195–205.
- 23 Gabbard GO: *Long-term psychodynamic psychotherapy*. Washington, American Psychiatric Publishing, 2004.
- 24 OPD Task Force: *Operationalized Psychodynamic Diagnosis OPD-2. Manual for Diagnosis and Treatment Planning*. Göttingen, Hogrefe & Huber, 2008.
- 25 Cohen J: *Statistical Power Analysis for the Behavioral Sciences*. Hillsdale, Lawrence Erlbaum, 1988.
- 26 Kazdin AE, Bass D: Power to detect differences between alternative treatments in comparative psychotherapy outcome research. *J Consult Clin Psychol* 1989;57:138–147.
- 27 Hofmann SG, Wu JQ, Boettcher H: Effect of cognitive-behavioral therapy for anxiety disorders on quality of life: a meta-analysis. *J Consult Clin Psychol* 2014;82:375–391.
- 28 Walker E, Nowacki AS: Understanding equivalence and noninferiority testing. *J Gen Intern Med* 2011;26:192–196.
- 29 Barker LE, Luman ET, McCauley MM, Chu SY: Assessing equivalence: an alternative to the use of difference tests for measuring disparities in vaccination coverage. *Am J Epidemiol* 2002;156:1056–1061.
- 30 Rogers JL, Howard KI, Vessey JT: Using significance tests to evaluate equivalence between two experimental groups. *Psychol Bull* 1993;113:553–565.
- 31 Lesaffre E: Superiority, equivalence, and non-inferiority trials. *Bull NYU Hosp Jt Dis* 2008;66:150–154.
- 32 Kraemer HC, Kupfer DJ: Size of treatment effects and their importance to clinical research and practice. *Biol Psychiatry* 2006;59:990–996.
- 33 Piaggio G, Elbourne DR, Pocock SJ, Evans SJ, Altman DG: Reporting of noninferiority and equivalence randomized trials: extension of the consort 2010 statement. *JAMA* 2012;308:2594–2604.
- 34 Malan DH: *The Frontier of Brief Psychotherapy*. New York, Plenum, 1976.
- 35 Sifneos PE: Short-term anxiety-provoking psychotherapy; in Davanloo H (ed): *Short-Term Dynamic Psychotherapy*. New York, Spectrum, 1978, pp 35–42.
- 36 Davanloo H: *Short-term Dynamic Psychotherapy*. New York, Jason Aronson, 1980.
- 37 Mann J: *Time-Limited Psychotherapy*. Cambridge, Harvard University Press, 1973.
- 38 Strupp HH, Binder JL: *Psychotherapy in a New Key: A Guide to Time-Limited Dynamic Psychotherapy*. New York, Basic Books, 1984.
- 39 Horowitz M, Kaltreider N: Brief therapy of the stress response syndrome. *Psychiatr Clin North Am* 1979;2:365–377.
- 40 Burnand Y, Andreoli A, Kolatte E, Venturini A, Rosset N: Psychodynamic psychotherapy and clomipramine in the treatment of major depression. *Psychiatr Serv* 2002;53:585–590.
- 41 de Jonghe F, Kool S, van Aalst G, Dekker J, Peen J: Combining psychotherapy and antidepressants in the treatment of depression. *J Affect Disord* 2001;64:217–229.
- 42 Thompson LW, Gallagher D, Breckenridge JS: Comparative effectiveness of psychotherapies for depressed elders. *J Consult Clin Psychol* 1987;55:385–390.
- 43 Milrod B, Leon AC, Busch F, Rudden M, Schwaberg M, Clarkin J, Aronson A, Singer M, Turchin W, Klass ET, Graf E, Teres JJ, Shear MK: A randomized controlled clinical trial of psychoanalytic psychotherapy for panic disorder. *Am J Psychiatry* 2007;164:265–272.
- 44 Clarkin JF, Levy KN, Lenzenweger MF, Kernberg OF: Evaluating three treatments for borderline personality disorder: a multiwave study. *Am J Psychiatry* 2007;164:922–928.
- 45 Woody GE, Luborsky L, McLellan AT, O'Brien CP, Beck AT, Blaine A, Herman I, Hole A: Psychotherapy for opiate addicts: does it help? *Arch Gen Psychiatry* 1983;40:639–645.
- 46 Woody GE, Luborsky L, McLellan AT, O'Brien CP: Psychotherapy in community methadone programs: a validation study. *Am J Psychiatry* 1995;152:1302–1308.
- 47 Wiborg IM, Dahl AA: Does brief dynamic psychotherapy reduce the relapse rate of panic disorder? *Arch Gen Psychiatry* 1996;53:689–694.
- 48 Winston A, Laikin M, Pollack J, Samstag LW, McCullough L, Muran JC: Short-term psychotherapy of personality disorders. *Am J Psychiatry* 1994;151:190–194.
- 49 de Jonghe F, Hendricksen M, van Aalst G, Kool S, Peen V, Van R, van den Eijnden E, Dekker J: Psychotherapy alone and combined with pharmacotherapy in the treatment of depression. *Br J Psychiatry* 2004;185:37–45.
- 50 Driessen E, Van HL, Don FJ, Peen J, Kool S, Westra D, Hendriksen M, Schoevers RA, Cuijpers P, Twisk J, Dekker JJ: The efficacy of cognitive-behavioural therapy and psychodynamic therapy in the outpatient treatment of major depression: a randomized clinical trial. *Am J Psychiatry* 2013;170:1041–1050.
- 51 Crits-Christoph P, Siqueland L, Blaine J, Frank A, Luborsky L, Onken LS, Muenz LR, Thase ME, Weiss RD, Gastfriend DR, Woody GE, Barber JP, Butler SF, Daley D, Salloum I, Bishop S, Najavits LM, Lis J, Mercer D, Griffin ML, Moras K, Beck AT: Psychosocial treatments for cocaine dependence: National Institute on Drug Abuse Collaborative Cocaine Treatment Study. *Arch Gen Psychiatry* 1999;56:493–502.
- 52 Leichsenring F, Salzer S, Beutel ME, Herpertz S, Hiller W, Hoyer J, Huesing J, Leibing E: Psychodynamic therapy and cognitive therapy in social anxiety disorder – a multi-center randomized controlled trial. *Am J Psychiatry* 2013;170:759–767.

- 53 Knekt P, Lindfors O, Harkanen T, Valikoski M, Virtala E, Laaksonen MA, Marttunen M, Kaipainen M, Renlund C: Randomized trial on the effectiveness of long- and short-term psychodynamic psychotherapy and solution-focused therapy on psychiatric symptoms during a 3-year follow-up. *Psychol Med* 2008; 38:689–703.
- 54 Bögels SM, Wijts P, Oort FJ, Sallaerts SJ: Psychodynamic psychotherapy versus cognitive behavior therapy for social anxiety disorder: an efficacy and partial effectiveness trial. *Depress Anxiety* 2014;31:363–373.
- 55 Zipfel S, Wild B, Gross G, Friederich H-C, Teufel M, Schellberg D, Herzog W: Focal psychodynamic therapy, cognitive behaviour therapy, and optimised treatment as usual in outpatients with anorexia nervosa (ANTOP study): randomised controlled trial. *Lancet* 2013;383:127–137.
- 56 Vinnars B, Barber JP, Noren K, Gallop R, Weinryb RM: Manualized supportive-expressive psychotherapy versus nonmanualized community-delivered psychodynamic therapy for patients with personality disorders: bridging efficacy and effectiveness. *Am J Psychiatry* 2005;162:1933–1940.
- 57 Hobson R: *Forms of Feeling. The Heart of Psychotherapy*. London, Tavistock, 1985.
- 58 Shapiro DA, Firth JA: *Exploratory Therapy Manual for the Sheffield Psychotherapy Project (SAPU Memo 733)*. Sheffield, University of Sheffield, 1985.
- 59 Cooper PJ, Murray L, Wilson A, Romaniuk H: Controlled trial of the short- and long-term effect of psychological treatment of post-partum depression. I. Impact on maternal mood. *Br J Psychiatry* 2003;182:412–419.
- 60 Johansson R, Björklund M, Hornborg C, Karlsson S, Hesser H, Ljotsson B, Rousseau A, Frederick JJ, Andersson G: Affect-focused psychodynamic psychotherapy for depression and anxiety through the internet: a randomized controlled trial. *PeerJ* 2013;102:1–22.
- 61 Johansson R, Ekbladh S, Hebert A: Psychodynamic guided self-help for adult depression through the internet: a randomised controlled trial. *PLoS One* 2012;7:e38021.
- 62 Maina G, Forner F, Bogetto F: Randomized controlled trial comparing brief dynamic and supportive therapy with waiting list condition in minor depressive disorders. *Psychother Psychosom* 2005;74:3–50.
- 63 Maina G, Rosso G, Crespi C, Bogetto F: Combined brief dynamic therapy and pharmacotherapy in the treatment of major depressive disorder: a pilot study. *Psychother Psychosom* 2007;76:298–305.
- 64 Beutel ME, Weissflog G, Leuteritz K, Wiltink J, Haselbacher A, Ruckes C, Kuhnt S, Barthel Y, Imruck BH, Zwerenz R, Brahler E: Efficacy of short-term psychodynamic psychotherapy (STPP) with depressed breast cancer patients: results of a randomized controlled multicenter trial. *Ann Oncol* 2014;25:378–384.
- 65 Thase ME: Comparative effectiveness of psychodynamic psychotherapy and cognitive-behavioral therapy: It's about time, and what's next? *Am J Psychiatry* 2013;170:953–955.
- 66 Salminen JK, Karlsson H, Hietala J, J K, Aalto S, Markkula J, Rasi-Hakala H, Toikka T: Short-term psychodynamic psychotherapy and fluoxetine in major depressive disorder: a randomized comparative study. *Psychother Psychosom* 2008;77:351–357.
- 67 Shapiro DA, Barkham M, Rees A, Hardy GE, Reynolds S, Startup M: Effects of treatment duration and severity of depression on the effectiveness of cognitive-behavioral and psychodynamic-interpersonal psychotherapy. *J Consult Clin Psychol* 1994;62:522–534.
- 68 Barber J, Barrett M, Gallop R, Rynn M, Rickels K: Short-term dynamic psychotherapy versus pharmacotherapy for major depressive disorder: a randomized, placebo-controlled trial. *J Clin Psychiatry* 2012;73:66–73.
- 69 McCallum M, Piper WE: A controlled study of effectiveness and patient suitability for short-term group psychotherapy. *Int J Group Psychother* 1990;40:431–452.
- 70 Piper WE, McCallum M, Joyce AS, Ogradniczuk J: Patient personality and time-limited group psychotherapy for complicated grief. *Int J Group Psychother* 2001;51:525–552.
- 71 Knijnik DZ, Kapczynski F, Chachamovich E, Margis R, Eizirik CL: Psychodynamic group treatment for generalized social phobia. *Rev Bras Psiquiatr* 2004;26:77–781.
- 72 Leichsenring F, Hoyer J, Beutel M, Herpertz S, Hiller W, Irie E, Joraschky P, König HH, de Liz TM, Nolting B, Pöhlmann K, Salzer S, Schauenburg H, Stangier U, Strauss B, Subic-Wrana C, Vormfelde S, Weniger G, Willutzki U, Wiltink J, Leibing E: The social phobia psychotherapy research network. The first multicenter randomized controlled trial of psychotherapy for social phobia: rationale, methods and patient characteristics. *Psychother Psychosom* 2009;78:35–41.
- 73 Leichsenring F, Salzer S, Beutel ME, Herpertz S, Hiller W, Hoyer J, Huesing J, Joraschky P, Nolting B, Poehlmann K, Ritter V, Stangier U, Strauss B, Tefikow S, Teismann T, Willutzki U, Wiltink J, Leibing E: Long-term outcome of psychodynamic therapy and cognitive-behavioral therapy in social anxiety disorder. *Am J Psychiatry* 2014;171:1074–1082.
- 74 Knijnik DZ, Blanco C, Salum GA, Moraes CU, Mombach C, Almeida E, Pereira M, Strapasson A, Manfro GG, Eizirik CL: A pilot study of clonazepam versus psychodynamic group therapy plus clonazepam in the treatment of generalized social anxiety disorder. *Eur Psychiatry* 2008;23:567–574.
- 75 Knijnik DZ, Salum GAJ, Blanco C, Moraes C, Hauck S, Mombach CK, Strapasson AC, Manfro GG, Eizirik CL: Defense style changes with the addition of psychodynamic group therapy to clonazepam in social anxiety disorder. *J Nerv Ment Dis* 2009;197:547–551.
- 76 Beutel ME, Scheurich V, Knebel A, Michal M, Wiltink J, Graf-Morgenstern M, Tschan R, Milrod B, Wellek S, Subic-Wrana C: Implementing panic-focused psychodynamic psychotherapy into clinical practice. *Can J Psychiatry* 2013;58:326–334.
- 77 Andersson G, Paxling B, Roch-Norlund P, Östman G, Norgren A, Almlöv J, Georén L, Breitholtz E, Dahlin M, Cuijpers P, Carlbring P, Silverberg F: Internet-based psychodynamic versus cognitive behavioral guided self-help for generalized anxiety disorder: a randomized controlled trial. *Psychother Psychosom* 2012;81:344–355.
- 78 Leichsenring F, Salzer S, Jager U, Kächele U, Kreische R, Leweke F, Rüger U, Winkelbach C, Leibing E: Short-term psychodynamic psychotherapy and cognitive-behavioral therapy in generalized anxiety disorder: a randomized controlled trial. *Am J Psychiatry* 2009;166:875–881.
- 79 Bressi C, Porcellana M, Marinaccio PM, Nocito EP, Magri L: Short-term psychodynamic psychotherapy versus treatment as usual for depressive and anxiety disorders: a randomized clinical trial of efficacy. *J Nerv Ment Dis* 2010;198:647–652.
- 80 Brom D, Kleber RJ, Defares PB: Brief psychotherapy for posttraumatic stress disorders. *J Consult Clin Psychol* 1989;57:607–612.
- 81 Maina G, Rosso G, Rigardetto S, Chiado Piat S, Bogetto F: No effect of adding brief dynamic therapy to pharmacotherapy in the treatment of obsessive-compulsive disorder with concurrent major depression. *Psychother Psychosom* 2010;79:295–302.
- 82 Monsen K, Monsen TJ: Chronic pain and psychodynamic body therapy. *Psychotherapy* 2000;37:257–269.
- 83 Sattel H, Lahmann C, Gündel H, Guthrie E, Kruse J, Noll-Hussong M, Ohmann C, Ronel J, Sack M, Sauer N, Schneider G, Henningsen P: Brief psychodynamic interpersonal psychotherapy for patients with multisomatoform disorder: randomised controlled trial. *Br J Psychiatry* 2012;200:60–67.
- 84 Bachar E, Latzer Y, Kreitler S, Berry EM: Empirical comparison of two psychological therapies. Self-psychology and cognitive orientation in the treatment of anorexia and bulimia. *J Psychother Pract Res* 1999;8:115–128.
- 85 Poulsen S, Lunn S, Daniel SI, Folke S, Mathiesen BB, Katznelson H, Fairburn CG: A randomized controlled trial of psychoanalytic psychotherapy or cognitive-behavioral therapy for bulimia nervosa. *Am J Psychiatry* 2014;171:109–116.
- 86 Tasca G, Hilsenroth M, Thompson-Brenner H: Psychoanalytic psychotherapy or cognitive-behavioral therapy for bulimia nervosa. *Am J Psychiatry* 2014;17:583–584.
- 87 Poulsen S, Lunn S: Response to Tasca et al. *Am J Psychiatry* 2014;171:584.
- 88 Garner DM, Rockert W, Davis R, Garner MV, Olmsted MP, Eagle M: Comparison of cognitive-behavioral and supportive-expressive therapy for bulimia nervosa. *Am J Psychiatry* 1993;150:37–46.
- 89 Fairburn CG, Kirk J, O'Connor M, Cooper PJ: A comparison of two psychological treatments for bulimia nervosa. *Behav Res Ther* 1986;24:629–643.
- 90 Gowers D, Norton K, Halek C, Vrisp AH: Outcome of outpatient psychotherapy in a random allocation treatment study of anorexia nervosa. *Int J Eat Disord* 1994;15:165–177.

- 91 Dare C, Eisler I, Russel G, Treasure J, Dodge L: Psychological therapies for adults with anorexia nervosa: randomised controlled trial of out-patient treatments. *Br J Psychiatry* 2001;178:216–221.
- 92 Tasca GA, Ritchie K, Conrad G, Balfour L, Gayrton J, Lybanon V, Bissada H: Attachment scales predict outcome in a randomized clinical trial of group psychotherapy for binge eating disorder: an aptitude by treatment interaction. *Psychother Res* 2006;16:106–121.
- 93 Woody GE, Luborsky L, McLellan AT, O'Brien CP: Corrections and revised analyses for psychotherapy in methadone maintenance patients. *Arch Gen Psychiatry* 1990;47:788–789.
- 94 Sandahl C, Herlitz K, Ahlin G, Rönning S: Time-limited group psychotherapy for moderately alcohol dependent patients: a randomized controlled clinical trial. *Psychother Res* 1998;8:361–378.
- 95 Levy KN, Meehan KB, Kelly KM, Reynoso JS, Weber M, Clarkin JF, Kernberg OF: Change in attachment patterns and reflective function in a randomized control trial of transference-focused psychotherapy for borderline personality disorder. *J Consult Clin Psychol* 2006;74:1027–1040.
- 96 Doering S, Horz S, Rentrop M, Fischer-Kern M, Schuster P, Benecke C, Buchheim A, Martius P, Buchheim P: Transference-focused psychotherapy v. treatment by community psychotherapists for borderline personality disorder: randomised controlled trial. *Br J Psychiatry* 2010;196:389–395.
- 97 Gregory RJ, Chlebowski S, Kang D, Remen AL, Soderberg MG, Stepkovitch J, Virk S: A controlled trial of psychodynamic psychotherapy for co-occurring borderline personality disorder and alcohol use disorder. *Psychotherapy* 2008;45:28–41.
- 98 Bateman A, Fonagy P: Randomized controlled trial of outpatient mentalization-based treatment versus structured clinical management for borderline personality disorder. *Am J Psychiatry* 2009;166:1355–1364.
- 99 Clarkin J, Kernberg OF, Yeomans F: *Transference-Focused Psychotherapy for Borderline Personality Disorder Patients*. New York, Guilford, 1999.
- 100 Munroe-Blum H, Marziali E: A controlled trial of short-term group treatment for borderline personality disorder. *J Personal Disord* 1995;9:190–198.
- 101 Giesen-Bloo J, van Dyck R, Spinhoven P, van Tilburg W, Dirksen C, van Asselt T, Kremers I, Nadort M, Arntz A: Outpatient psychotherapy for borderline personality disorder: randomized trial of schema-focused therapy vs transference-focused psychotherapy. *Arch Gen Psychiatry* 2006;63:649–658.
- 102 Yeomans F: Questions concerning the randomized trial of schema-focused therapy vs transference-focused psychotherapy. *Arch Gen Psychiatry* 2007;64:610–611.
- 103 Giesen-Bloo J, Arntz A: Questions concerning the randomized trial of schema-focused therapy vs transference-focused psychotherapy – reply. *Arch Gen Psychiatry* 2007;64:610–611.
- 104 Svartberg M, Stiles T, Seltzer MH: Randomized, controlled trial of the effectiveness of short-term dynamic psychotherapy and cognitive therapy for cluster C personality disorders. *Am J Psychiatry* 2004;161:810–817.
- 105 Muran JC, Safran JD, Samstag LW, Winston A: Evaluating an alliance-focused treatment for personality disorders. *Psychotherapy* 2005;42:532–545.
- 106 Emmelkamp P, Benner A, Kuipers A, Feiertag G, Koster HC, van Appeldorn FJ: Comparison of brief dynamic and cognitive-behavioral therapies in avoidant personality disorder. *Br J Psychiatry* 2006;189:60–64.
- 107 Hellerstein DJ, Rosenthal RN, Pinsker H, Samstag LW, Muran JC, Winston A: A randomized prospective study comparing supportive and dynamic therapies. Outcome and alliance. *J Psychother Pract Res* 1998;7:261–271.
- 108 Abbass AA, Sheldon A, Gyra J, Kalpin A: Intensive short-term psychotherapy for DSM-IV personality disorders: a randomized controlled trial. *J Nerv Ment Dis* 2008;196:211–216.
- 109 Guthrie E, Kapur N, Mackway-Jones K, Chew-Graham C, Moorey J, Mendel E, Marino-Francis F, Sanderson S, Turpin C, Boddy G, Tomenson B: Randomised controlled trial of brief psychological intervention after deliberate self-poisoning. *BMJ* 2001;323:135–138.
- 110 Guthrie E, Moorey J, Margison F, Barker H, Palmer S, McGrath G, Tomenson B, Creed F: Cost-effectiveness of brief psychodynamic-interpersonal therapy in high utilizers of psychiatric services. *Arch Gen Psychiatry* 1999;56:519–526.
- 111 Rosenbaum B, Harder S, Knudsen P, Koster A, Lajer M, Lindhardt A, Valbak K, Winther G: Supportive psychodynamic psychotherapy versus treatment as usual for first-episode psychosis: two-year outcome. *Psychiatry* 2012;75:331–341.
- 112 Snyder DK, Wills RM: Behavioral vs. insight-oriented marital therapy: effects on individual and interspousal functioning. *J Consult Clin Psychol* 1989;57:39–46.
- 113 Snyder DK, Wills RM, Grady-Fletcher A: Long-term effectiveness of behavioral versus insight-oriented marital therapy: a 4-year follow-up study. *J Consult Clin Psychol* 1991;59:138–141.
- 114 Lazar SG, Yeomans FE: Guest editors' introduction. *Psychodyn Psychiatry* 2014;42:347–352.
- 115 Parloff MB: Psychotherapy research evidence and reimbursement decisions: Bambi meets Godzilla. *Am J Psychiatry* 1982;139:718–727.
- 116 Glass RM: Psychodynamic psychotherapy and research evidence: Bambi survives Godzilla? *JAMA* 2008;300:1587–1589.
- 117 de Jonghe F: Psychoanalytic supportive psychotherapy. *J Am Psychoanal Assoc* 1994;42:421–446.
- 118 Abbass AA, Kisely S, Kroenke K: Short-term psychodynamic psychotherapy for somatic disorders. Systematic review and meta-analysis of clinical trials. *Psychother Psychosom* 2009;265–274.
- 119 Richardson WS, Doster LM: Comorbidity and multimorbidity need to be placed in the context of a framework of risk, responsiveness, and vulnerability. *J Clin Epidemiol* 2014;67:244–246.
- 120 Berk M, Parker G: The elephant on the couch: side effects of psychotherapy. *Aust NZ J Psychiatry* 2009;43:787–794.
- 121 Greenson RR: *The Techniques and Practice of Psychoanalysis*. New York, International Universities Press, 1967.
- 122 Abbass A, Sheldon A, Gyra J, Kalpin A: Intensive short-term dynamic psychotherapy for DSM-IV personality disorders: a randomized controlled trial. *J Nerv Ment Dis* 2008;196:211–216.
- 123 Cuijpers P, Sijbrandij M, Koole SL, Andersson G, Beekman AT, Reynolds CF 3rd: The efficacy of psychotherapy and pharmacotherapy in treating depressive and anxiety disorders: a meta-analysis of direct comparisons. *World Psychiatry* 2013;12:137–148.
- 124 Keefe JR, McCarthy KS, Dinger U, Zilchamano S, Barber JP: A meta-analytic review of psychodynamic therapies for anxiety disorders. *Clin Psychol Rev* 2014;34:309–323.
- 125 Cuijpers P, Berking M, Andersson G, Quigley L, Kleiboer A, Dobson KS: A meta-analysis of cognitive-behavioural therapy for adult depression, alone and in comparison with other treatments. *Can J Psychiatry* 2013;58:376–385.
- 126 Leichsenring F, Salzer S, Hilsenroth M, Leibling E, Leweke F, Rabung S: Treatment integrity: an unresolved issue in psychotherapy research. *Curr Psychiatry Rev* 2011;7:313–321.
- 127 Baardseth TP, Goldberg SB, Pace BT, Wislocki AP, Frost ND, Siddiqui JR, Lindemann AM, Kivlighan DM 3rd, Laska KM, Del Re AC, Minami T, Wampold BE: Cognitive-behavioral therapy versus other therapies: redux. *Clin Psychol Rev* 2013;33:395–405.
- 128 Gallagher-Thompson DE, Steffen AM: Comparative effects of cognitive-behavioral and brief psychodynamic psychotherapies for depressed family caregivers. *J Consult Clin Psychol* 1994;62:543–549.
- 129 Hollon SD, DeRubeis RJ, Shelton RC, Amsterdam JD, Salomon RM, O'Reardon JP, Lovett ML, Young PR, Haman KL, Freeman BB, Gallop R: Prevention of relapse following cognitive therapy vs medications in moderate to severe depression. *Arch Gen Psychiatry* 2005;62:417–422.
- 130 Shea MT, Elkin I, Imber SD, Sotsky SM, Watkins JT, Collins JF, Pilkonis PA, Beckham E, Glass DR, Dolan RT, et al: Course of depressive symptoms over follow-up. Findings from the National Institute of Mental Health Treatment of Depression Collaborative Research Program. *Arch Gen Psychiatry* 1992;49:782–787.

- 131 Gallagher-Thompson DE, Hanley-Peterson P, Thompson LW: Maintenance of gains versus relapse following brief psychotherapy for depression. *J Consult Clin Psychol* 1990;58:371–374.
- 132 Ablon JS, Jones EE: Validity of controlled clinical trials of psychotherapy: findings from the NIMH Treatment of Depression Collaborative Research Program. *Am J Psychiatry* 2002;159:775–783.
- 133 Ablon JS, Jones EE: How expert clinicians prototypes of an ideal treatment correlate with outcome in psychodynamic and cognitive-behavioral therapy. *Psychother Res* 1998;8:71–83.
- 134 Ablon JS, Levy KN, Katzenstein T: Beyond brand names of psychotherapy: identifying empirically supported change processes. *Psychotherapy (Chic)* 2006;43:216–231.
- 135 Luborsky L, Woody GE, McLellan AT, Rosenzweig J: Can independent judges recognize different psychotherapies? An experiment with manual-guided therapies. *J Consult Clin Psychol* 1982;30:49–62.
- 136 Crits-Christoph P, Connolly Gibbons MB, Mukherjee D: Psychotherapy process-outcome research; in Lambert MJ (ed): *Bergin and Garfield's Handbook of Psychotherapy and Behavior Change*. New York, Wiley, 2013.
- 137 Barber JP, Crits-Christoph P, Luborsky L: Effects of therapist adherence and competence on patient outcome in brief dynamic therapy. *J Consult Clin Psychol* 1996;64:619–622.
- 138 Hilsenroth M, Ackerman SJ, Blagys MD, Baitty MR, Mooney MA: Short-term psychodynamic psychotherapy for depression: an examination of statistical, clinically significant, and technique-specific change. *J Nerv Ment Dis* 2003;191:349–357.
- 139 Slavin-Mulford J, Hilsenroth M, Weinberger J, Gold J: Therapeutic interventions related to outcome in psychodynamic psychotherapy for anxiety disorder patients. *J Nerv Ment Dis* 2011;199:214–221.
- 140 Pitman S, Slavin-Mulford J, Hilsenroth M: Psychodynamic techniques related to outcome for anxiety disorder patients at different points in treatment. *J Nerv Ment Dis* 2014;202:391–396.
- 141 Luborsky L, McLellan AT, Woody GE, O'Brien CP, Auerbach A: Therapy success and its determinants. *Arch Gen Psychiatry* 1985;42:602–611.
- 142 Crits-Christoph P, Luborsky L: Changes in CCRT pervasiveness during psychotherapy; in Luborsky L, Crits-Christoph P (eds): *Understanding Transference: The CCRT Method*. New York, Basic Books, 1990, pp 133–146.
- 143 Connolly Gibbons MB, Crits-Christoph P, Barber JP, Stirman SW, Gallop R, Goldstein LA, Temes CM, Ring-Kurtz S: Unique and common mechanisms of change across cognitive and dynamic psychotherapies. *J Consult Clin Psychol* 2009;77:801–813.
- 144 Norton PJ, Phillip LM: Transdiagnostic approaches to the treatment of anxiety disorders: a quantitative review. *Psychotherapy (Chic)* 2008;45:214–226.
- 145 Leichsenring F, Salzer S: A unified protocol for the transdiagnostic psychodynamic treatment of anxiety disorders: an evidence-based approach. *Psychotherapy (Chic)* 2014;51:224–245.
- 146 Leichsenring F, Schauenburg H: Empirically supported methods of short-term psychodynamic therapy in depression – towards an evidence-based unified protocol. *J Affect Disord* 2014;169C:128–143.
- 147 Busch FN, Singer MB, Milrod BL, Aronson AC: *Manual of Panic-Focused Psychodynamic Psychotherapy – Extended Range*. New York, Taylor & Francis, 2012.
- 148 Thoma NC, McKay D, Gerber AJ, Milrod BL, Edwards AR, Kocsis JH: A quality-based review of randomized controlled trials of cognitive-behavioral therapy for depression: an assessment and metaregression. *Am J Psychiatry* 2012;169:22–30.
- 149 Edwards SJ, Lilford RJ, Braunholtz D, Jackson J: Why 'underpowered' trials are not necessarily unethical. *Lancet* 1997;350:804–807.
- 150 Schulz KF, Grimes DA: Sample size calculations in randomised trials: mandatory and mystical. *Lancet* 2005;365:1348–1353.
- 151 Borenstein M, Hedges LV, Higgins JPT, Rothstein HR: *Introduction to Meta-Analysis*. Chichester, Wiley, 2011.
- 152 Abbass A, Rabung S, Leichsenring F, Refseth J, Midgeley N: Psychodynamic psychotherapy for children and adolescents: a meta-analysis of short-term psychodynamic model. *J Am Child Adolesc Psychiatry* 2013;52:863–875.
- 153 Kernberg OF: A psychoanalytic model for the classification of personality disorders; in Achenheil M, Bondy B, Engel R, Ermann M, Nedopil N (eds): *Implications of Psychopharmacology to Psychiatry*. New York, Springer, 1996.
- 154 Cramer B, Robert-Tissot C, Stern D: Outcome evaluation in brief mother-infant psychotherapy: a preliminary report. *Infant Ment Health J* 1990;2:278–300.
- 155 Stern D: *The Motherhood Constellation*. New York, Basic Books, 1995.
- 156 de Jonghe F, de Maat S, Van R, Hendriksen M, Kool S, van Aalst G, Dekker J: Short-term psychoanalytic supportive psychotherapy for depressed patients. *Psychoanal Inquiry* 2013;33:614–625.
- 157 Silverberg F: *Make the Leap: A Practical Guide to Breaking the Patterns That Hold You Back*. New York, Marlowe, 2005.
- 158 Malan DH: *A Study of Brief Psychotherapy*. New York, Plenum, 1963.
- 159 Haselbacher A, Barthel Y, Brähler E, Imruck B, Kuhnt S, Zwerenz R, Beutel ME: Psychodynamic short-term therapy for cancer patients suffering from depression. A treatment manual on the basis of Luborsky's supportive-expressive therapy (in German). *Psychotherapeut* 2010;55:321–328.
- 160 Malan DH, Osimo F (eds): *Psychodynamics, Training and Outcome in Brief Psychotherapy*. Oxford, Butterworth-Heinemann, 1992.
- 161 McCullough L, Kuhn N, Andrews S, Kaplan A, Wolf J, Hurley CL: *Treating Affect Phobia. A manual for Short-Term Dynamic Psychotherapy*. New York, Guilford Press, 2003.
- 162 Frederick RJ: *Living Like You Mean It: Use the Wisdom and Power of Your Emotions to Get the Life You Really Want*. San Francisco, Jossey-Bass, 2009.
- 163 McCallum M, Piper W: Psychoanalytically oriented short-term groups for outpatients; unsettled issues. *Group* 1988;12:21–32.
- 164 McCallum M, Piper W, Joyce AS: *Manual for time-limited, short-term, supportive group therapy for patients experiencing pathological bereavement*. Unpubl Manuscript, 1995.
- 165 Leichsenring F, Beutel ME, Leibing E: Psychodynamic psychotherapy for social phobia: a treatment manual based on supportive-expressive therapy. *Bull Menninger Clin* 2007;71:56–83.
- 166 Milrod B, Busch F, Cooper A, Shapiro T: *Manual of Panic-Focused Psychodynamic Psychotherapy*. Washington, American Psychiatric Press, 1997.
- 167 Monsen K, Monsen TJ: Affects and affect consciousness. A psychotherapy model of integrating Silvan Tomkins' affect and script theory within the framework of self-psychology; in Goldberg A (ed): *Pluralism in Self-Psychology: Progress in Self-Psychology*. Hillsdale, Analytic Press, 1999, vol 15, pp 287–306.
- 168 Hardy G, Barkham M, Shapiro D, Guthrie E, Margison F: *Psychodynamic-Interpersonal Psychotherapy*. New York, Sage, 2011.
- 169 PISO Working Group: *Psychodynamic-Interpersonal Therapy for Somatoform Disorders (PISO) – A Manualized Short-Term Psychotherapy*. Göttingen, Hogrefe, 2011.
- 170 Bachar E: The contribution of self-psychology to the treatment of anorexia and bulimia. *Am J Psychother* 1998;52:147–165.
- 171 Dare C: Psychoanalytic psychotherapy (of eating disorders); in Gabbard GO (ed): *Treatment of Psychiatric Disorders*. Washington, American Psychiatric Press, 1995, pp 2129–2151.
- 172 Schauenburg H, Friederich HC, Wild B, Zipfel S, Herzog W: *Focal psychodynamic psychotherapy of anorexia nervosa: a treatment manual*. *Psychotherapeut* 2009;54:270–280.
- 173 Bateman A, Fonagy P: Mentalization-based treatment of BPD. *J Personal Disord* 2004;18:36–51.
- 174 Gregory RJ, Remen AL: A manual-based psychodynamic therapy for treatment-resistant borderline personality disorder. *Psychotherapy (Chic)* 2008;45:15–27.
- 175 Guthrie E, Moorey J, Barker H, Margison F, McGrath G: Psychodynamic interpersonal psychotherapy in patients with treatment resistant psychiatric symptoms. *Br J Psychother* 1998;115:155–166.
- 176 Nadelson CC, Paolino TJ: Marital therapy from a psychoanalytic perspective; in Paolino TJ, Crady BS (eds): *Marriage and Marital Therapy: Psychoanalytic, Behavioral and Systems Theory Perspectives*. New York, Brunner-Mazel, 1978, pp 89–164.